arXiv:1201.5473v1 [math.RT] 26 Jan 2012

A New Factor from E_6-Mod to E_7-Mod [1]

Xiaoping Xu

Hua Loo-Keng Key Mathematical Laboratory

Institute of Mathematics, Academy of Mathematics & System Sciences

Chinese Academy of Sciences, Beijing 100190, P.R. China [2]

Abstract

We find a new representation of the simple Lie algebra of type E_7 on the polynomial algebra in 27 variables, which gives a fractional representation of the corresponding Lie group on 27-dimensional space. Using this representation and Shen's idea of mixed product, we construct a functor from E_6-**Mod** to E_7-**Mod**. A condition for the functor to map a finite-dimensional irreducible E_6-module to an infinite-dimensional irreducible E_7-module is obtained. Our general frame also gives a direct polynomial extension from irreducible E_6-modules to irreducible E_7-modules. The obtained infinite-dimensional irreducible E_7-modules are $(\mathcal{G}, K)$-modules in terms of Lie group representations. The results could be used in studying the quantum field theory with E_7 symmetry and symmetry of partial differential equations.

1 Introduction

A quantum field is an operator-valued function on a certain Hilbert space, which is often a direct sum of infinite-dimensional irreducible modules of a certain Lie algebra (group). The Lie algebra of two-dimensional conformal group is exactly the Virasoro algebra, which contains a central element. The minimal models of two-dimensional conformal field theory were constructed from direct sums of certain infinite-dimensional irreducible modules of the Virasoro algebra, where the value taken by the central element is an important physical quantity, called the *central charge*.

It is well known that n-dimensional projective group gives rise to a non-homogenous representation of the Lie algebra $sl(n+1,\mathbb{C})$ on the polynomial functions of the projective space. Using Shen's mixed product for Witt algebras, Zhao and the author [ZX] generalized the above representation of $sl(n+1,\mathbb{C})$ to a non-homogenous representation on the tensor space of any finite-dimensional irreducible $gl(n,\mathbb{C})$-module with the polynomial space. Moreover, the structure of such a representation was completely determined. The result can be used to study the quantum field theory with $sl(n+1,\mathbb{C})$ as the symmetry. Furthermore, we [XZ] generalize the conformal representation of $o(n+2,\mathbb{C})$ to a non-homogenous representation of $o(n+2,\mathbb{C})$ on the tensor space of any finite-dimensional irreducible $o(n,\mathbb{C})$-module with a polynomial space by Shen's idea of mixed product for Witt algebras. It turns out that a hidden central transformation is involved. More importantly, we find a condition on the constant value taken by the central transformation such that the generalized conformal representation is irreducible. The result would be useful in higher-dimensional conformal field theory.

---

[1] 2000 Mathematical Subject Classification. Primary 17B10, 17B25; Secondary 17B01.

[2] Research supported by China NSF 10871193

In [X4], we find a new representation of the simple Lie algebra of type E_6 on the polynomial algebra in 16 variables, which gives a fractional representation of the corresponding Lie group on 16-dimensional space. Using this representation and Shen's idea of mixed product, we construct a functor from D_5-**Mod** to E_6-**Mod**. A condition for the functor to map a finite-dimensional irreducible D_5-module to an infinite-dimensional irreducible E_6-module is obtained. Our general frame also gives a direct polynomial extension from irreducible D_5-modules to irreducible E_6-modules. The obtained infinite-dimensional irreducible E_6-modules are $(\mathcal{G}, K)$-modules in terms of Lie group representations. The results could be used in studying the quantum field theory with E_6 symmetry and symmetry of partial differential equations.

This paper is the forth work in the program of studying quantum-field motivated representations of finite-dimensional simple Lie algebras. It is well known that the minimal dimension of irreducible modules over the simple Lie algebra of type E_7 is 56. Based on a grading of the simple Lie algebra of type E_7, we find a first-order differential operator representation of the Lie algebra on the polynomial algebra in 27 independent variables. In fact, the corresponding Lie group representation is given by fractional transformations on 27-dimensional space. Using this representation and Shen's idea of mixed product, we construct a new functor from E_6-**Mod** to E_7-**Mod**, where a hidden central transformation is involved. More importantly, a condition for the functor to map a finite-dimensional irreducible E_6-module to an infinite-dimensional irreducible E_7-module in terms of the constant value taken by the central transformation is obtained. The well-known Dickson cubic invariant (cf. [Dl]) plays an important role in our approach. Our general frame also gives a direct polynomial extension from irreducible E_6-modules to irreducible E_7-modules, which can be applied to obtain explicit bases of irreducible E_7-modules from those of irreducible E_6-modules. The result could be useful in understanding the quantum field theory with E_7 symmetry. Our fractional representation of the E_7 Lie group could also be used in symmetry analysis of partial differential equations just as the conformal representation of orthogonal Lie groups does. Our infinite-dimensional irreducible E_7-modules are $(\mathcal{G}, K)$-modules in terms of the corresponding Lie group representations.

The E_7 Lie algebra and group are important mathematical objects with broad applications. They are second most complicated finite-dimensional simple Lie algebra and group. The complexity indeed imply rich connotation. Here we are only able to list a small part of it. Ramond [R] (1977) gave a group theoretical analysis of a symmetry breaking affected by Higgs fields for the vector-like unified theory base on E_7. Cvitanović [Co] (1981) studied the E_7 symmetry with negative bosonic dimension. Han, Kim and Tanii [HKT] (1986) presented a supersymmetrization of the six-dimensional anomaly-free $E_6 \times E_7 \times U(1)$ theory with Lorentz Chern-Simons term. Kato and Kitazawa [KK] (1989) studied the correlation functions in an E_7-type modular-invariant Wess-Zumino-Witten theory, which is related to the scheme of string compatification proposed by Gepner. Ma [M] (1990) found the spectrum-dependent solutions to the Yang-Baxter equation for quantum E_7.

Fernádez, Garcia Fuerres and Perelomov [FFP] (2005) re-expressed the quantum Calogero-Sutherland model for the Lie algebra E_7 and the particular value of the coupling constant $k = 1$, using the fundamental irreducible characters of the algebra as dynamic variables. Moreover, they used the model to obtain explicitly the characters and Clebsch-

Gordan series for the algebra E_7. D'Auria, Ferrara and Trigiante [DFT] (2006) considered M-theory compactified on seven-torus with fluxes when all the seven anti-symmetric tensor fields in four dimensions have been dualized into scalars and thus the $E_{7(7)}$ symmetry was recovered. Duff and Ferrara [DF] (2007) proposed that a particular tripartite entanglement of seven quits, encoded in the Fano plane, is described by the E_7 group and that the entanglement measure is given by Cartan's quartic E_7 invariant. Brink, Kim and Ramond [BKR] (2008) used the Cremmer-Julia $E_{7(7)}$ nonlinear symmetry of $N = 8$ supergravity to derive its order κ on-shell Hamltonian in terms of the chiral light-cone superfield. Borsten [Bl] (2008) used the Freudenthal triple system to sketch the precise dictionary relating the 56 charges, parametrizing the general black hole solution, to the 56-dimensional quantum states of E_7.

Part of the mathematical story of E_7 is as follows. Brown [Br1] (1968) constructed a central simple nonassociative algebraic structure on the Freudenthal triple system of E_7 and obtained a new interpretation of the E_7 Lie algebra. He [Br2] (1969) proved that the group leaving Cartan's quartic form invariant modulo its center is exactly a simple E_7 Lie group. Faulkner [Fau] (1972) studied the geometry of the planes of the octave symplectic geometry of the ternary algebra related to the minimal representation of E_7. Ferrar [Fer] (1980) used the Freudenthal triple system to classified the simple E_7 Lie algebras over algebraic number fields. Kleidman and Ryba [KR] (1993) proved a Kostant's conjecture in the case of E_7, which says that the group $PSL(2, 37)$ embeds in an E_7 Lie group. Moreover, Griess and Ryba [GR] (1994) proved that the group $U_3(8)$ embeds in an E_7 Lie group and the group $Sz(8)$ does not. Ginzburg [Gd] (1995) showed that the twisted partial L-function on the 56-dimensional representation of $GE_7(\mathbb{C})$ is entire except the points 0 and 1.

Cooperstein [Cb] (1995) proved that Cartan's quartic E_7-invariant is the unique fundamental invariant over the basic module. Moreover, Shult [Se] (1997) showed that the basic representation of E_7 provides the absolutely universal embedding of the point-line geometry $E_{7,1}$. Plotkin [P] (1998) investigated the stability of the K_1-functor the E_7 group. Doković [Dd1] (1999) classified nilpotent adjoint orbits of real simple noncompact groups of type E_7 by means of Caley triples. He [Dd2] (2001) also worked out the partial order on nilpotent orbits in the split real Lie algebra of type E_7. The E_7 root system was used by Sekiguchi [Sj] (1999) to study the configurations of seven lines on the real projective plane. Kleidman, Meierfrankenfeld and Ryba [KMR] (1999) proved that the group HS embeds in $E_7(5)$. Choi and Yoon [CY] (1999) calculated the homology of the double and triple loop spaces of E_7.

Ukai [U] (2001) computed the b-functions of the prehomogeneous vector space arising from a cuspidal character sheaf of E_7. Garibaidi [Gr] (2001) gave explicit descriptions of the homogeneous projective varieties associated with the E_7 group with trivial Tits algebras. A characterization of quadratic forms of type E_7 was given by De Medits [Dt] (2002). Cherenousov [Cv] (2003) used Hasse principle to prove that the Rost invariant has trivial kernel for a quasi-split E_7 group. Kono, Lin and Nishimura [KLN] (2003) characterized the mod 3 cohomology of E_7. Weiss [W] (2006) solved a fundamental question about the structure of the automorphism group of the Moufang quadrangles of type E_7. Moreover, we found in [X3] that the weight matrices of E_7 on its minimal irreducible module and adjoint module generate ternary orthogonal codes with large minimal distances.

This work further reveals new beauties of the simple Lie algebra of type E_7. In Section 2, we construct the 27-dimensional representation of the simple Lie algebra of type E_6 in terms of first-order differential operators on the polynomial algebra in 27 independent variables from the lattice-construction of the simple Lie algebra of type E_7. In Section 3, we realize the simple Lie algebra of type E_7 in terms of first-order differential operators on the polynomial algebra in 27 independent variables. Section 4 is devoted to the explicit presentation of the functor from E_6-**Mod** to E_7-**Mod**. Finally in Section 6, we determine a condition for the functor to map a finite-dimensional irreducible E_6-module to an infinite-dimensional irreducible E_7-module.

2 Polynomial Representation of E_6 via E_7

In this section, we will explicitly construct the 27-dimensional basic irreducible representation of E_6.

For convenience, we will use the notion

$$\overline{i, i+j} = \{i, i+1, i+2, ..., i+j\} \tag{2.1}$$

for integer i and positive integer j throughout this paper. We start with the root lattice construction of the simple Lie algebra of type E_7. As we all known, the Dynkin diagram of E_7 is as follows:



Let $\{\alpha_i \mid i \in \overline{1,7}\}$ be the simple positive roots corresponding to the vertices in the diagram, and let Φ_{E_7} be the root system of E_7. Set

$$Q_{E_7} = \sum_{i=1}^{7} \mathbb{Z}\alpha_i, \tag{2.2}$$

the root lattice of type E_7. Denote by $(\cdot,\cdot)$ the symmetric $\mathbb{Z}$-bilinear form on Q_{E_7} such that

$$\Phi_{E_7} = \{\alpha \in Q_{E_7} \mid (\alpha,\alpha) = 2\}. \tag{2.3}$$

Define $F(\cdot,\cdot) : Q_{E_7} \times Q_{E_7} \to \{\pm 1\}$ by

$$F(\sum_{i=1}^{7} k_i\alpha_i, \sum_{j=1}^{7} l_j\alpha_j) = (-1)^{\sum_{i=1}^{7} k_i l_i + \sum_{1\leq i<j\leq 7} k_i l_j (\alpha_i,\alpha_j)}, \qquad k_i, l_j \in \mathbb{Z}. \tag{2.4}$$

Then for $\alpha,\beta,\gamma \in Q_{E_7}$,

$$F(\alpha+\beta,\gamma) = F(\alpha,\gamma)F(\beta,\gamma), \;\; F(\alpha,\beta+\gamma) = F(\alpha,\beta)F(\alpha,\gamma), \tag{2.5}$$

$$F(\alpha,\beta)F(\beta,\alpha)^{-1} = (-1)^{(\alpha,\beta)}, \;\; F(\alpha,\alpha) = (-1)^{(\alpha,\alpha)/2}. \tag{2.6}$$

In particular,

$$F(\alpha,\beta) = -F(\beta,\alpha) \qquad \text{if } \alpha,\beta,\alpha+\beta \in \Phi_{E_7}. \tag{2.7}$$

Denote

$$H_{E_7}=\sum_{i=1}^{7}\mathbb{R}\alpha_i. \tag{2.8}$$

The simple Lie algebra of type E_7 is

$$\mathcal{G}^{E_7}=H_{E_7}\oplus\bigoplus_{\alpha\in\Phi_{E_7}}\mathbb{R}E_\alpha \tag{2.9}$$

with the Lie bracket $[\cdot,\cdot]$ determined by:

$$[H_{E_7},H_{E_7}]=0,\;\; [h,E_\alpha]=(h,\alpha)E_\alpha,\;\; [E_\alpha,E_{-\alpha}]=-\alpha, \tag{2.10}$$

$$[E_\alpha,E_\beta]=\begin{cases}0 & \text{if } \alpha+\beta\not\in\Phi_{E_7},\\ F(\alpha,\beta)E_{\alpha+\beta} & \text{if } \alpha+\beta\in\Phi_{E_7}\end{cases} \tag{2.11}$$

for $\alpha,\beta\in\Phi_{E_7}$ and $h\in H_{E_7}$ (e.g., cf. [K, X1]). Moreover, we define a bilinear form $(\cdot|\cdot)$ on $\mathcal{G}^{E_6}$ by

$$(h_1|h_2)=(h_1,h_2),\;\; (h|E_\alpha)=0,\;\; (E_\alpha|E_\beta)=-\delta_{\alpha+\beta,0} \tag{2.12}$$

for $h_1,h_2\in H$ and $\alpha,\beta\in\Phi_{E_7}$. It can be verified that $(\cdot|\cdot)$ is a $\mathcal{G}^{E_7}$-invariant form, that is,

$$([u,v]|w)=-(v|[u,w])\qquad \text{for }\; u,v\in\mathcal{G}^{E_7}. \tag{2.13}$$

Note that

$$Q_{E_6}=\sum_{i=1}^{6}\mathbb{Z}\alpha_i\subset Q_{E_7} \tag{2.14}$$

is the root lattice of E_6 and

$$\Phi_{E_6}=Q_{E_6}\bigcap\Phi_{E_7} \tag{2.15}$$

is the root system of E_6. Set

$$H_{E_6}=\sum_{i=1}^{6}\mathbb{R}\alpha_i. \tag{2.16}$$

Then the subalgebra

$$\mathcal{G}^{E_6}=H_{E_6}\oplus\bigoplus_{\alpha\in\Phi_{E_6}}\mathbb{R}E_\alpha \tag{2.17}$$

of $\mathcal{G}^{E_7}$ is exactly the simple Lie algebra of type E_6. Denote by $\Phi_{E_6}^+$ the set of positive roots of E_6 and by $\Phi_{E_7}^+$ the set of positive roots of E_7. The elements of $\Phi_{E_6}^+$ are:

$$\alpha_1+2\alpha_2+2\alpha_3+3\alpha_4+2\alpha_5+\alpha_6, \tag{2.18}$$

$$\{\alpha_1+\sum_{r=3}^{j}\alpha_r\mid j\in\overline{2,6}\}\bigcup\{\sum_{r=i+1}^{j}\alpha_r\mid 2\leq i<j\leq 6\}, \tag{2.19}$$

$$\{\sum_{s=2}^{j}\alpha_s+\sum_{t=4}^{k}\alpha_t\mid 2\leq j<k\leq 6\} \tag{2.20}$$

and

$$\{\sum_{\iota=1}^{i}\alpha_\iota+\sum_{s=3}^{j}\alpha_s+\sum_{t=4}^{k}\alpha_t\mid 2\leq i<j<k\leq 6\}. \tag{2.21}$$

Denote by $\bar{\Phi}_{E_7}^+$ the set of the following positive roots:

$$\alpha_1+\sum_{r=3}^{7}\alpha_r,\qquad \alpha_3+2\alpha_4+\alpha_5+\sum_{i=1}^{6}\alpha_i+\sum_{r=1}^{7}\alpha_r, \tag{2.22}$$

$$\{2\sum_{s=1}^{6}\alpha_s-\alpha_1+\alpha_4-\alpha_6+\sum_{r=i+1}^{7}\alpha_r \mid i\in\overline{1,6}\},\qquad \{\sum_{r=i+1}^{7}\alpha_r \mid i\in\overline{2,6}\}, \tag{2.23}$$

$$\{\sum_{s=2}^{j}\alpha_s+\sum_{t=4}^{7}\alpha_t \mid j\in\overline{2,6}\},\qquad \{\sum_{\iota=1}^{i}\alpha_\iota+\sum_{s=3}^{j}\alpha_s+\sum_{t=4}^{7}\alpha_t \mid 2\leq i<j\leq 6\}. \tag{2.24}$$

Then

$$\Phi_{E_7}^+=\Phi_{E_6}^+\bigcup\bar{\Phi}_{E_7}^+. \tag{2.25}$$

For convenience, we also denote

$$E_{(k_1,\dots,k_r)}=E_\alpha,\;\; E'_{(k_1,\dots,k_r)}=E_{-\alpha}\qquad \text{for }\;\alpha=\sum_{s=1}^{r}k_s\alpha_s\in\Phi_{E_7}^+,\; k_r\neq 0. \tag{2.26}$$

Write

$$\xi_1=E_{(0,0,0,0,0,0,1)},\;\;\xi_2=E_{(0,0,0,0,0,1,1)},\;\;\xi_3=E_{(0,0,0,0,1,1,1)},\;\;\xi_4=E_{(0,0,0,1,1,1,1)}, \tag{2.27}$$

$$\xi_5=E_{(0,0,1,1,1,1,1)},\;\;\xi_6=E_{(0,1,0,1,1,1,1)},\;\;\xi_7=E_{(0,1,1,1,1,1,1)},\;\;\xi_8=E_{(1,0,1,1,1,1,1)}, \tag{2.28}$$

$$\xi_9=E_{(0,1,1,2,1,1,1)},\;\;\xi_{10}=E_{(1,1,1,1,1,1,1)},\;\;\xi_{11}=E_{(0,1,1,2,2,1,1)},\;\;\xi_{12}=E_{(1,1,1,2,1,1,1)}, \tag{2.29}$$

$$\xi_{13}=E_{(1,1,1,2,2,1,1)},\;\;\xi_{14}=E_{(0,1,1,2,2,2,1)},\;\;\xi_{15}=E_{(1,1,2,2,1,1,1)},\;\;\xi_{16}=E_{(1,1,2,2,2,1,1)}, \tag{2.30}$$

$$\xi_{17}=E_{(1,1,1,2,2,2,1)},\;\;\xi_{18}=E_{(1,1,2,3,2,1,1)},\;\;\xi_{19}=E_{(1,1,2,2,2,2,1)},\;\;\xi_{20}=E_{(1,2,2,3,2,1,1)}, \tag{2.31}$$

$$\xi_{21}=E_{(1,1,2,3,2,2,1)},\;\;\xi_{22}=E_{(1,1,2,3,3,2,1)},\;\;\xi_{23}=E_{(1,2,2,3,2,2,1)},\;\;\xi_{24}=E_{(1,2,2,3,3,2,1)}, \tag{2.32}$$

$$\xi_{25}=E_{(1,2,2,4,3,2,1)},\;\;\xi_{26}=E_{(1,2,3,4,3,2,1)},\;\;\xi_{27}=E_{(2,2,3,4,3,2,1)}, \tag{2.33}$$

$$\eta_1=E'_{(0,0,0,0,0,0,1)},\;\;\eta_2=E'_{(0,0,0,0,0,1,1)},\;\;\eta_3=E'_{(0,0,0,0,1,1,1)},\;\;\eta_4=E'_{(0,0,0,1,1,1,1)}, \tag{2.34}$$

$$\eta_5=E'_{(0,0,1,1,1,1,1)},\;\;\eta_6=E'_{(0,1,0,1,1,1,1)},\;\;\eta_7=E'_{(0,1,1,1,1,1,1)},\;\;\eta_8=E'_{(1,0,1,1,1,1,1)}, \tag{2.35}$$

$$\eta_9=E'_{(0,1,1,2,1,1,1)},\;\;\eta_{10}=E'_{(1,1,1,1,1,1,1)},\;\;\eta_{11}=E'_{(0,1,1,2,2,1,1)},\;\;\eta_{12}=E'_{(1,1,1,2,1,1,1)}, \tag{2.36}$$

$$\eta_{13}=E'_{(1,1,1,2,2,1,1)},\;\;\eta_{14}=E'_{(0,1,1,2,2,2,1)},\;\;\eta_{15}=E'_{(1,1,2,2,1,1,1)},\;\;\eta_{16}=E'_{(1,1,2,2,2,1,1)}, \tag{2.37}$$

$$\eta_{17}=E'_{(1,1,1,2,2,2,1)},\;\;\eta_{18}=E'_{(1,1,2,3,2,1,1)},\;\;\eta_{19}=E'_{(1,1,2,2,2,2,1)},\;\;\eta_{20}=E'_{(1,2,2,3,2,1,1)}, \tag{2.38}$$

$$\eta_{21}=E'_{(1,1,2,3,2,2,1)},\;\;\eta_{22}=E'_{(1,1,2,3,3,2,1)},\;\;\eta_{23}=E'_{(1,2,2,3,2,2,1)},\;\;\eta_{24}=E'_{(1,2,2,3,3,2,1)}, \tag{2.39}$$

$$\eta_{25}=E'_{(1,2,2,4,3,2,1)},\;\;\eta_{26}=E'_{(1,2,3,4,3,2,1)},\;\;\eta_{27}=E'_{(2,2,3,4,3,2,1)}. \tag{2.40}$$

Set

$$\mathcal{G}_-=\sum_{i=1}^{27}\mathbb{C}\eta_i,\qquad \mathcal{G}_0=\mathcal{G}^{E_6}+\mathbb{C}\alpha_7,\qquad \mathcal{G}_+=\sum_{i=1}^{27}\mathbb{C}\eta_i. \tag{2.41}$$

Then $\mathcal{G}_\pm$ are abelian subalgebras of $\mathcal{G}^{E_7}$ and $\mathcal{G}_0$ is a maximal reductive Lie subalgebra of $\mathcal{G}^{E_7}$. Moreover,

$$[\mathcal{G}_+,\mathcal{G}_-]\subset\mathcal{G}_0,\qquad [\mathcal{G}_0,\mathcal{G}_\pm]\subset\mathcal{G}_\pm,\qquad \mathcal{G}^{E_7}=\mathcal{G}_-\oplus\mathcal{G}_0\oplus\mathcal{G}_+. \tag{2.42}$$

Denote by λ_i the ith fundamental weight of $\mathcal{G}^{E_6}$. With respect to the adjoint representation of $\mathcal{G}^{E_7}$, $\mathcal{G}_+$ forms an irreducible $\mathcal{G}^{E_6}$-module with highest weight λ_1 and $\mathcal{G}_-$ forms an irreducible $\mathcal{G}^{E_6}$-module with highest weight λ_6.

Set

$$\mathcal{A} = \mathbb{C}[x_1, x_2, ..., x_{27}], \tag{2.43}$$

the polynomial algebra in $x_1, x_2, ..., x_{27}$. Write

$$[u, \eta_i] = \sum_{j=1}^{27} \varphi_{i,j}(u)\eta_j \qquad \text{for } \; i \in \overline{1,27}, \; u \in \mathcal{G}_0, \tag{2.44}$$

where $\varphi_{i,j}(u) \in \mathbb{C}$. Define an action of $\mathcal{G}_0$ on $\mathcal{A}$ by

$$u(f) = \sum_{i,j=1}^{27} \varphi_{i,j}(u) x_j \partial_{x_i}(f) \qquad \text{for } \; u \in \mathcal{G}_0, \; f \in \mathcal{A}. \tag{2.45}$$

Then $\mathcal{A}$ forms a $\mathcal{G}_0$-module and the subspace

$$V = \sum_{i=1}^{27} \mathbb{C}x_i \tag{2.46}$$

forms a $\mathcal{G}_0$-submodule isomorphic to $\mathcal{G}_-$, where the isomorphism is determined by $x_i \mapsto \eta_i$ for $i \in \overline{1,27}$.

Denote by $\mathbb{N}$ the set of nonnegative integers. Write

$$x^\alpha = \prod_{i=1}^{27} x_i^{\alpha_i}, \;\; \partial^\alpha = \prod_{i=1}^{27} \partial_{x_i}^{\alpha_i} \qquad \text{for } \; \alpha = (\alpha_1, \alpha_2, ..., \alpha_{27}) \in \mathbb{N}^{27}. \tag{2.47}$$

Let

$$\mathbb{A} = \sum_{\alpha \in \mathbb{N}^{27}} \mathcal{A}\partial^\alpha \tag{2.48}$$

be the algebra of differential operators on $\mathcal{A}$. Then the linear transformation τ determined by

$$\tau(x^\beta \partial^\gamma) = x^\gamma \partial^\beta \qquad \text{for } \; \beta, \gamma \in \mathbb{N}^{27} \tag{2.49}$$

is an involutive anti-automorphism of $\mathbb{A}$.

Thanks to (2.10), (2.11), (2.44) and (2.45), we find the following representation formulas of $\mathcal{G}_0$:

$$E_{\alpha_1}|_{\mathcal{A}} = x_5\partial_{x_8} + x_7\partial_{x_{10}} + x_9\partial_{x_{12}} + x_{11}\partial_{x_{13}} + x_{14}\partial_{x_{17}} + x_{26}\partial_{x_{27}}, \tag{2.50}$$

$$E_{\alpha_2}|_{\mathcal{A}} = x_4\partial_{x_6} + x_5\partial_{x_7} + x_8\partial_{x_{10}} - x_{18}\partial_{x_{20}} - x_{21}\partial_{x_{23}} - x_{22}\partial_{x_{24}}, \tag{2.51}$$

$$E_{\alpha_3}|_{\mathcal{A}} = x_4\partial_{x_5} + x_6\partial_{x_7} + x_{12}\partial_{x_{15}} + x_{13}\partial_{x_{16}} + x_{17}\partial_{x_{19}} - x_{25}\partial_{x_{26}}, \tag{2.52}$$

$$E_{\alpha_4}|_{\mathcal{A}} = x_3\partial_{x_4} - x_7\partial_{x_9} - x_{10}\partial_{x_{12}} - x_{16}\partial_{x_{18}} - x_{19}\partial_{x_{21}} - x_{24}\partial_{x_{25}}, \tag{2.53}$$

$$E_{\alpha_5}|_{\mathcal{A}} = x_2\partial_{x_3} - x_9\partial_{x_{11}} - x_{12}\partial_{x_{13}} - x_{15}\partial_{x_{16}} - x_{21}\partial_{x_{22}} - x_{23}\partial_{x_{24}}, \tag{2.54}$$

$$E_{\alpha_6}|_{\mathcal{A}} = x_1\partial_{x_2} - x_{11}\partial_{x_{14}} - x_{13}\partial_{x_{17}} - x_{16}\partial_{x_{19}} - x_{18}\partial_{x_{21}} - x_{20}\partial_{x_{23}}, \tag{2.55}$$

$$E_{(1,0,1)}|_{\mathcal{A}} = x_4\partial_{x_8} + x_6\partial_{x_{10}} - x_9\partial_{x_{15}} + x_{11}\partial_{x_{16}} - x_{14}\partial_{x_{19}} + x_{25}\partial_{x_{27}}, \tag{2.56}$$

$$E_{(0,1,0,1)}|_{\mathcal{A}} = x_3\partial_{x_6} + x_5\partial_{x_9} + x_8\partial_{x_{12}} + x_{16}\partial_{x_{20}} + x_{19}\partial_{x_{23}} - x_{22}\partial_{x_{25}}, \tag{2.57}$$

$$E_{(0,0,1,1)}|_{\mathcal{A}} = x_3\partial_{x_5} + x_6\partial_{x_9} - x_{10}\partial_{x_{15}} + x_{13}\partial_{x_{18}} + x_{17}\partial_{x_{21}} + x_{24}\partial_{x_{26}}, \tag{2.58}$$

$$E_{(0,0,0,1,1)}|_{\mathcal{A}} = x_2\partial_{x_4} - x_7\partial_{x_{11}} - x_{10}\partial_{x_{13}} + x_{15}\partial_{x_{18}} - x_{19}\partial_{x_{22}} + x_{23}\partial_{x_{25}}, \tag{2.59}$$

$$E_{(0,0,0,0,1,1)}|_{\mathcal{A}} = x_1\partial_{x_3} - x_9\partial_{x_{14}} - x_{12}\partial_{x_{17}} - x_{15}\partial_{x_{19}} + x_{18}\partial_{x_{22}} + x_{20}\partial_{x_{24}}, \tag{2.60}$$

$$E_{(1,0,1,1)}|_{\mathcal{A}} = x_3\partial_{x_8} + x_6\partial_{x_{12}} + x_7\partial_{x_{15}} - x_{11}\partial_{x_{18}} - x_{14}\partial_{x_{21}} + x_{24}\partial_{x_{27}}, \tag{2.61}$$

$$E_{(0,1,1,1)}|_{\mathcal{A}} = x_3\partial_{x_7} - x_4\partial_{x_9} + x_8\partial_{x_{15}} - x_{13}\partial_{x_{20}} - x_{17}\partial_{x_{23}} + x_{22}\partial_{x_{26}}, \tag{2.62}$$

$$E_{(0,1,0,1,1)}|_{\mathcal{A}} = x_2\partial_{x_6} + x_5\partial_{x_{11}} + x_8\partial_{x_{13}} - x_{15}\partial_{x_{20}} + x_{19}\partial_{x_{x_{24}}} + x_{21}\partial_{x_{25}}, \tag{2.63}$$

$$E_{(0,0,1,1,1)}|_{\mathcal{A}} = x_2\partial_{x_5} + x_6\partial_{x_{11}} - x_{10}\partial_{x_{16}} - x_{12}\partial_{x_{18}} + x_{17}\partial_{x_{22}} - x_{23}\partial_{x_{26}}, \tag{2.64}$$

$$E_{(0,0,0,1,1,1)}|_{\mathcal{A}} = x_1\partial_{x_4} - x_7\partial_{x_{14}} - x_{10}\partial_{x_{17}} + x_{15}\partial_{x_{21}} + x_{16}\partial_{x_{22}} - x_{20}\partial_{x_{25}}, \tag{2.65}$$

$$E_{(1,1,1,1)}|_{\mathcal{A}} = x_3\partial_{x_{10}} - x_4\partial_{x_{12}} - x_5\partial_{x_{15}} + x_{11}\partial_{x_{20}} + x_{14}\partial_{x_{23}} + x_{22}\partial_{x_{27}}, \tag{2.66}$$

$$E_{(1,0,1,1,1)}|_{\mathcal{A}} = x_2\partial_{x_8} + x_6\partial_{x_{13}} + x_7\partial_{x_{16}} + x_9\partial_{x_{18}} - x_{14}\partial_{x_{22}} - x_{23}\partial_{x_{27}}, \tag{2.67}$$

$$E_{(0,1,1,1,1)}|_{\mathcal{A}} = x_2\partial_{x_7} - x_4\partial_{x_{11}} + x_8\partial_{x_{16}} + x_{12}\partial_{x_{20}} - x_{17}\partial_{x_{24}} - x_{21}\partial_{x_{26}}, \tag{2.68}$$

$$E_{(0,1,0,1,1,1)}|_{\mathcal{A}} = x_1\partial_{x_6} + x_5\partial_{x_{14}} + x_8\partial_{x_{17}} - x_{15}\partial_{x_{23}} - x_{16}\partial_{x_{24}} - x_{18}\partial_{x_{25}}, \tag{2.69}$$

$$E_{(0,0,1,1,1,1)}|_{\mathcal{A}} = x_1\partial_{x_5} + x_6\partial_{x_{14}} - x_{10}\partial_{x_{19}} - x_{12}\partial_{x_{21}} - x_{13}\partial_{x_{22}} + x_{20}\partial_{x_{26}}, \tag{2.70}$$

$$E_{(1,1,1,1,1}|_{\mathcal{A}} = x_2\partial_{x_{10}} - x_4\partial_{x_{13}} - x_5\partial_{x_{16}} - x_9\partial_{x_{20}} + x_{14}\partial_{x_{24}} - x_{21}\partial_{x_{27}}, \tag{2.71}$$

$$E_{(1,0,1,1,1,1)}|_{\mathcal{A}} = x_1\partial_{x_8} + x_6\partial_{x_{17}} + x_7\partial_{x_{19}} + x_9\partial_{x_{21}} + x_{11}\partial_{x_{22}} + x_{20}\partial_{x_{27}}, \tag{2.72}$$

$$E_{(0,1,1,2,1)}|_{\mathcal{A}} = x_2\partial_{x_9} - x_3\partial_{x_{11}} + x_8\partial_{x_{18}} - x_{10}\partial_{x_{20}} - x_{17}\partial_{x_{25}} + x_{19}\partial_{x_{26}}, \tag{2.73}$$

$$E_{(0,1,1,1,1,1)}|_{\mathcal{A}} = x_1\partial_{x_7} - x_4\partial_{x_{14}} + x_8\partial_{x_{19}} + x_{12}\partial_{x_{23}} + x_{13}\partial_{x_{24}} + x_{18}\partial_{x_{26}}, \tag{2.74}$$

$$E_{(1,1,1,2,1)}|_{\mathcal{A}} = x_2\partial_{x_{12}} - x_3\partial_{x_{13}} - x_5\partial_{x_{18}} + x_7\partial_{x_{20}} + x_{14}\partial_{x_{25}} + x_{19}\partial_{x_{27}}, \tag{2.75}$$

$$E_{(1,1,1,1,1,1)}|_{\mathcal{A}} = x_1\partial_{x_{10}} - x_4\partial_{x_{17}} - x_5\partial_{x_{19}} - x_9\partial_{x_{23}} - x_{11}\partial_{x_{24}} + x_{18}\partial_{x_{27}}, \tag{2.76}$$

$$E_{(0,1,1,2,1,1)}|_{\mathcal{A}} = x_1\partial_{x_9} - x_3\partial_{x_{14}} + x_8\partial_{x_{21}} - x_{10}\partial_{x_{23}} + x_{13}\partial_{x_{25}} - x_{16}\partial_{x_{26}}, \tag{2.77}$$

$$E_{(1,1,2,2,1)}|_{\mathcal{A}} = x_2\partial_{x_{15}} - x_3\partial_{x_{16}} + x_4\partial_{x_{18}} - x_6\partial_{x_{20}} - x_{14}\partial_{x_{26}} - x_{17}\partial_{x_{27}}, \tag{2.78}$$

$$E_{(1,1,1,2,1,1)}|_{\mathcal{A}} = x_1\partial_{x_{12}} - x_3\partial_{x_{17}} - x_5\partial_{x_{21}} + x_7\partial_{x_{23}} - x_{11}\partial_{x_{25}} - x_{16}\partial_{x_{27}}, \tag{2.79}$$

$$E_{(0,1,1,2,2,1)}|_{\mathcal{A}} = x_1\partial_{x_{11}} - x_2\partial_{x_{14}} + x_8\partial_{x_{22}} - x_{10}\partial_{x_{24}} - x_{12}\partial_{x_{25}} + x_{15}\partial_{x_{26}}, \tag{2.80}$$

$$E_{(1,1,2,2,1,1)}|_{\mathcal{A}} = x_1\partial_{x_{15}} - x_3\partial_{x_{19}} + x_4\partial_{x_{21}} - x_6\partial_{x_{23}} + x_{11}\partial_{x_{26}} + x_{13}\partial_{x_{27}}, \tag{2.81}$$

$$E_{(1,1,1,2,2,1)}|_{\mathcal{A}} = x_1\partial_{x_{13}} - x_2\partial_{x_{17}} - x_5\partial_{x_{22}} + x_7\partial_{x_{24}} + x_9\partial_{x_{25}} + x_{15}\partial_{x_{27}}, \tag{2.82}$$

$$E_{(1,1,2,2,2,1)}|_{\mathcal{A}} = x_1\partial_{x_{16}} - x_2\partial_{x_{19}} + x_4\partial_{x_{22}} - x_6\partial_{x_{24}} - x_9\partial_{x_{26}} - x_{12}\partial_{x_{27}}, \tag{2.83}$$

$$E_{(1,1,2,3,2,1)}|_{\mathcal{A}} = x_1\partial_{x_{18}} - x_2\partial_{x_{21}} + x_3\partial_{x_{22}} - x_6\partial_{x_{25}} + x_7\partial_{x_{26}} + x_{10}\partial_{x_{27}}, \tag{2.84}$$

$$E_{(1,2,2,3,2,1)}|_{\mathcal{A}} = x_1\partial_{x_{20}} - x_2\partial_{x_{23}} + x_3\partial_{x_{24}} - x_4\partial_{x_{25}} + x_5\partial_{x_{26}} + x_8\partial_{x_{27}}, \tag{2.85}$$

$$E_{-\alpha}|_{\mathcal{A}} = -\tau(E_\alpha|_{\mathcal{A}}) \qquad \text{for } \alpha \in \Phi^+_{E_6}, \tag{2.86}$$

$$\alpha_r|_{\mathcal{A}} = \sum_{i=1}^{27} a_{i,r} x_i \partial_{x_i} \qquad \text{for } r \in \overline{1,7} \tag{2.87}$$

with $a_{i,r}$ given in the following table:

Table 1

i	$a_{i,1}$	$a_{i,2}$	$a_{i,3}$	$a_{i,4}$	$a_{i,5}$	$a_{i,6}$	$a_{i,7}$	i	$a_{i,1}$	$a_{i,2}$	$a_{i,3}$	$a_{i,4}$	$a_{i,5}$	$a_{i,6}$	$a_{i,7}$
1	0	0	0	0	0	1	-2	2	0	0	0	0	1	-1	-1
3	0	0	0	1	-1	0	-1	4	0	1	1	-1	0	0	-1
5	1	1	-1	0	0	0	-1	6	0	-1	1	0	0	0	-1
7	1	-1	-1	1	0	0	-1	8	-1	1	0	0	0	0	-1
9	1	0	0	-1	1	0	-1	10	-1	-1	0	1	0	0	-1
11	1	0	0	0	-1	1	-1	12	-1	0	1	-1	1	0	-1
13	-1	0	1	0	-1	1	-1	14	1	0	0	0	0	-1	0
15	0	0	-1	0	1	0	-1	16	0	0	-1	1	-1	1	-1
17	-1	0	1	0	0	-1	0	18	0	1	0	-1	0	1	-1
19	0	0	-1	1	0	-1	0	20	0	-1	0	0	0	1	-1
21	0	1	0	-1	1	-1	0	22	0	1	0	0	-1	0	0
23	0	-1	0	0	1	-1	0	24	0	-1	0	1	-1	0	0
25	0	0	1	-1	0	0	0	26	1	0	-1	0	0	0	0
27	-1	0	0	0	0	0	0								

Let

$$\widehat{\alpha}=2\alpha_1+3\alpha_2+4\alpha_3+6\alpha_4+5\alpha_5+4\alpha_6+3\alpha_7. \tag{2.88}$$

Then

$$(\widehat{\alpha},\alpha_r)=0 \qquad \text{for } \; r\in\overline{1,6} \tag{2.89}$$

by the Dynkin diagram of E_7. Thanks to (2.10),

$$[\widehat{\alpha},\mathcal{G}^{D_6}]=0. \tag{2.90}$$

By Schur's Lemma, $\widehat{\alpha}|_V=c\sum_{i=1}^{27}x_i\partial_{x_i}$. According to the coefficients of $x_1\partial_{x_1}$ in (2.87) and Table 1, we have that

$$\widehat{\alpha}|_{\mathcal{A}}=\sum_{i=1}^{16}x_i\partial_{x_i}=D \tag{2.91}$$

is the degree operator on $\mathcal{A}$.

3 Extended Representation of E_7

In this section, we realize the simple Lie algebra of type E_7 in terms of first-order differential operators on the polynomial algebra in 27 independent variables.

Recall that a singular vector of $\mathcal{G}^{E_6}$ is a nonzero weight vector annihilated by positive root vectors. According to Table 1, we find a $\mathcal{G}^{E_6}$-singular vector

$$\zeta_1=x_1x_{14}+x_2x_{11}+x_3x_9+x_4x_7-x_5x_6 \tag{3.1}$$

of weight λ_1 as (3.1)-(3.6) in [X2]. Note

$$E_{-\alpha_1}|_{\mathcal{A}}=-x_8\partial_{x_5}-x_{10}\partial_{x_7}-x_{12}\partial_{x_9}-x_{13}\partial_{x_{11}}-x_{17}\partial_{x_{14}}-x_{27}\partial_{x_{26}}, \tag{3.2}$$

$$E_{-\alpha_2}|_{\mathcal{A}}=-x_6\partial_{x_4}-x_7\partial_{x_5}-x_{10}\partial_{x_8}+x_{20}\partial_{x_{18}}+x_{23}\partial_{x_{21}}+x_{24}\partial_{x_{22}}, \tag{3.3}$$

$$E_{-\alpha_3}|_{\mathcal{A}}=-x_5\partial_{x_4}-x_7\partial_{x_6}-x_{15}\partial_{x_{12}}-x_{16}\partial_{x_{13}}-x_{19}\partial_{x_{17}}+x_{26}\partial_{x_{25}}, \tag{3.4}$$

$$E_{-\alpha_4}|_{\mathcal{A}}=-x_4\partial_{x_3}+x_9\partial_{x_7}+x_{12}\partial_{x_{10}}+x_{18}\partial_{x_{16}}+x_{21}\partial_{x_{19}}+x_{25}\partial_{x_{24}}, \tag{3.5}$$

$$E_{-\alpha_5}|_{\mathcal{A}} = -x_3\partial_{x_2} + x_{11}\partial_{x_9} + x_{13}\partial_{x_{12}} + x_{16}\partial_{x_{15}} + x_{22}\partial_{x_{21}} + x_{24}\partial_{x_{23}}, \tag{3.6}$$

$$E_{-\alpha_6}|_{\mathcal{A}} = -x_2\partial_{x_1} + x_{14}\partial_{x_{11}} + x_{17}\partial_{x_{13}} + x_{19}\partial_{x_{16}} + x_{21}\partial_{x_{18}} + x_{23}\partial_{x_{20}} \tag{3.7}$$

by (2.50)-(2.55) and (2.86).

As (3.7)-(3.24) in [X2], we set by (3,2)-(3.7) that

$$\zeta_2 = x_1x_{17} + x_2x_{13} + x_3x_{12} + x_4x_{10} - x_6x_8, \tag{3.8}$$

$$\zeta_3 = x_1x_{19} + x_2x_{16} + x_3x_{15} + x_5x_{10} - x_7x_8, \tag{3.9}$$

$$\zeta_4 = -x_1x_{21} - x_2x_{18} + x_4x_{15} - x_5x_{12} + x_8x_9, \tag{3.10}$$

$$\zeta_5 = x_1x_{22} - x_3x_{18} - x_4x_{16} + x_5x_{13} - x_8x_{11}, \tag{3.11}$$

$$\zeta_6 = x_1x_{23} + x_2x_{20} + x_6x_{15} - x_7x_{12} + x_9x_{10}, \tag{3.12}$$

$$\zeta_7 = -x_1x_{24} + x_3x_{20} - x_6x_{16} + x_7x_{13} - x_{10}x_{11}, \tag{3.13}$$

$$\zeta_8 = x_2x_{22} + x_3x_{21} + x_4x_{19} - x_5x_{17} + x_8x_{14}, \tag{3.14}$$

$$\zeta_9 = x_1x_{25} + x_4x_{20} + x_6x_{18} - x_9x_{13} + x_{11}x_{12}, \tag{3.15}$$

$$\zeta_{10} = -x_2x_{24} - x_3x_{23} + x_6x_{19} - x_7x_{17} + x_{10}x_{14}, \tag{3.16}$$

$$\zeta_{11} = x_1x_{26} - x_5x_{20} - x_7x_{18} + x_9x_{16} - x_{11}x_{15}, \tag{3.17}$$

$$\zeta_{12} = x_2x_{25} - x_4x_{23} - x_6x_{21} + x_9x_{17} - x_{12}x_{14}, \tag{3.18}$$

$$\zeta_{13} = x_3x_{25} + x_4x_{24} + x_6x_{22} - x_{11}x_{17} + x_{13}x_{14}, \tag{3.19}$$

$$\zeta_{14} = x_1x_{27} - x_8x_{20} - x_{10}x_{18} + x_{12}x_{16} - x_{13}x_{15}, \tag{3.20}$$

$$\zeta_{15} = x_2x_{26} + x_5x_{23} + x_7x_{21} - x_9x_{19} + x_{14}x_{15}, \tag{3.21}$$

$$\zeta_{16} = x_3x_{26} - x_5x_{24} - x_7x_{22} + x_{11}x_{19} - x_{14}x_{16}, \tag{3.22}$$

$$\zeta_{17} = -x_2x_{27} - x_8x_{23} - x_{10}x_{21} + x_{12}x_{19} - x_{15}x_{17}, \tag{3.23}$$

$$\zeta_{18} = x_4x_{26} + x_5x_{25} + x_9x_{22} - x_{11}x_{21} + x_{14}x_{18}, \tag{3.24}$$

$$\zeta_{19} = -x_3x_{27} + x_8x_{24} + x_{10}x_{22} - x_{13}x_{19} + x_{16}x_{17}, \tag{3.25}$$

$$\zeta_{20} = -x_6x_{26} - x_7x_{25} + x_9x_{24} - x_{11}x_{23} + x_{14}x_{20}, \tag{3.26}$$

$$\zeta_{21} = -x_4x_{27} - x_8x_{25} - x_{12}x_{22} + x_{13}x_{21} - x_{17}x_{18}, \tag{3.27}$$

$$\zeta_{22} = x_5x_{27} - x_8x_{26} + x_{15}x_{22} - x_{16}x_{21} + x_{18}x_{19}, \tag{3.28}$$

$$\zeta_{23} = x_6x_{27} + x_{10}x_{25} - x_{12}x_{24} + x_{13}x_{23} - x_{17}x_{20}, \tag{3.29}$$

$$\zeta_{24} = -x_7x_{27} + x_{10}x_{26} + x_{15}x_{24} - x_{16}x_{23} + x_{19}x_{20}, \tag{3.30}$$

$$\zeta_{25} = -x_9x_{27} + x_{12}x_{26} + x_{15}x_{25} - x_{18}x_{23} + x_{20}x_{21}, \tag{3.31}$$

$$\zeta_{26} = -x_{11}x_{27} + x_{13}x_{26} + x_{16}x_{25} - x_{18}x_{24} + x_{20}x_{22}, \tag{3.32}$$

$$\zeta_{27} = -x_{14}x_{27} + x_{17}x_{26} + x_{19}x_{25} - x_{21}x_{24} + x_{22}x_{23}. \tag{3.33}$$

Then

$$U = \sum_{i=1}^{27} \mathbb{C}\zeta_i \tag{3.34}$$

forms an irreducible $\mathcal{G}^{E_6}$-module with highest weight λ_1.

By (2.50)-(2.55) and (3.1)-(3.33), we calculate

$$E_{\alpha_1}|_U = \zeta_1\partial_{\zeta_2} + \zeta_{11}\partial_{\zeta_{14}} - \zeta_{15}\partial_{\zeta_{17}} - \zeta_{16}\partial_{\zeta_{19}} - \zeta_{18}\partial_{\zeta_{21}} - \zeta_{20}\partial_{\zeta_{23}}, \tag{3.35}$$

$$E_{\alpha_2}|_U = \zeta_4\partial_{\zeta_6} + \zeta_5\partial_{\zeta_7} + \zeta_8\partial_{\zeta_{10}} - \zeta_{18}\partial_{\zeta_{20}} - \zeta_{21}\partial_{\zeta_{23}} - \zeta_{22}\partial_{\zeta_{24}}, \tag{3.36}$$

$$E_{\alpha_3}|_U = \zeta_2\partial_{\zeta_3} - \zeta_9\partial_{\zeta_{11}} - \zeta_{12}\partial_{\zeta_{15}} - \zeta_{13}\partial_{\zeta_{16}} - \zeta_{21}\partial_{\zeta_{22}} - \zeta_{23}\partial_{\zeta_{24}}, \tag{3.37}$$

$$E_{\alpha_4}|_U = \zeta_3\partial_{\zeta_4} + \zeta_7\partial_{\zeta_9} + \zeta_{10}\partial_{\zeta_{12}} + \zeta_{16}\partial_{\zeta_{18}} + \zeta_{19}\partial_{\zeta_{21}} - \zeta_{24}\partial_{\zeta_{25}}, \tag{3.38}$$

$$E_{\alpha_5}|_U = \zeta_4\partial_{\zeta_5} + \zeta_6\partial_{\zeta_7} + \zeta_{12}\partial_{\zeta_{13}} + \zeta_{15}\partial_{\zeta_{16}} + \zeta_{17}\partial_{\zeta_{19}} - \zeta_{25}\partial_{\zeta_{26}}, \tag{3.39}$$

$$E_{\alpha_6}|_U = \zeta_5\partial_{\zeta_8} + \zeta_7\partial_{\zeta_{10}} + \zeta_9\partial_{\zeta_{12}} + \zeta_{11}\partial_{\zeta_{15}} - \zeta_{14}\partial_{\zeta_{17}} - \zeta_{26}\partial_{\zeta_{27}}, \tag{3.40}$$

$$E_{-\alpha_1}|_U = -\zeta_2\partial_{\zeta_1} - \zeta_{14}\partial_{\zeta_{11}} + \zeta_{17}\partial_{\zeta_{15}} + \zeta_{19}\partial_{\zeta_{16}} + \zeta_{21}\partial_{\zeta_{18}} + \zeta_{23}\partial_{\zeta_{20}}, \tag{3.41}$$

$$E_{-\alpha_2}|_U = -\zeta_6\partial_{\zeta_4} - \zeta_7\partial_{\zeta_5} - \zeta_{10}\partial_{\zeta_8} + \zeta_{20}\partial_{\zeta_{18}} + \zeta_{23}\partial_{\zeta_{21}} + \zeta_{24}\partial_{\zeta_{22}}, \tag{3.42}$$

$$E_{-\alpha_3}|_U = -\zeta_3\partial_{\zeta_2} + \zeta_{11}\partial_{\zeta_9} + \zeta_{15}\partial_{\zeta_{12}} + \zeta_{16}\partial_{\zeta_{13}} + \zeta_{22}\partial_{\zeta_{21}} + \zeta_{24}\partial_{\zeta_{23}}, \tag{3.43}$$

$$E_{-\alpha_4}|_U = -\zeta_4\partial_{\zeta_3} - \zeta_9\partial_{\zeta_7} - \zeta_{12}\partial_{\zeta_{10}} - \zeta_{18}\partial_{\zeta_{16}} - \zeta_{21}\partial_{\zeta_{19}} + \zeta_{25}\partial_{\zeta_{24}}, \tag{3.44}$$

$$E_{-\alpha_5}|_U = -\zeta_5\partial_{\zeta_4} - \zeta_7\partial_{\zeta_6} - \zeta_{13}\partial_{\zeta_{12}} - \zeta_{16}\partial_{\zeta_{15}} - \zeta_{19}\partial_{\zeta_{17}} + \zeta_{26}\partial_{\zeta_{25}}, \tag{3.45}$$

$$E_{-\alpha_6}|_U = -\zeta_8\partial_{\zeta_5} - \zeta_{10}\partial_{\zeta_7} - \zeta_{12}\partial_{\zeta_9} - \zeta_{15}\partial_{\zeta_{11}} + \zeta_{17}\partial_{\zeta_{14}} + \zeta_{27}\partial_{\zeta_{26}}. \tag{3.46}$$

Moreover, (2.84) and Table 1 give

$$\alpha_r|_U = \sum_{i=1}^{27} b_{i,r}\zeta_i\partial_{\zeta_i} \qquad \text{for } r \in \overline{1,6} \tag{3.47}$$

with $b_{i,r}$ given in the following table:

Table 2

i	$b_{i,1}$	$b_{i,2}$	$b_{i,3}$	$b_{i,4}$	$b_{i,5}$	$b_{i,6}$	i	$b_{i,1}$	$b_{i,2}$	$b_{i,3}$	$b_{i,4}$	$b_{i,5}$	$b_{i,6}$
1	1	0	0	0	0	0	2	-1	0	1	0	0	0
3	0	0	-1	1	0	0	4	0	1	0	-1	1	0
5	0	1	0	0	-1	1	6	0	-1	0	0	1	0
7	0	-1	0	1	-1	1	8	0	1	0	0	0	-1
9	0	0	1	-1	0	1	10	0	-1	0	1	0	-1
11	1	0	-1	0	0	1	12	0	0	1	-1	1	-1
13	0	0	1	0	-1	0	14	-1	0	0	0	0	1
15	1	0	-1	0	1	-1	16	1	0	-1	1	-1	0
17	-1	0	0	0	1	-1	18	1	1	0	-1	0	0
19	-1	0	0	1	-1	0	20	1	-1	0	0	0	0
21	-1	1	1	-1	0	0	22	0	1	-1	0	0	0
23	-1	-1	1	0	0	0	24	0	-1	-1	1	0	0
25	0	0	0	-1	1	0	26	0	0	0	0	-1	1
27	0	0	0	0	0	-1							

Recall that the Lie bracket in the algebra $\mathbb{A}$ of differential operators (cf. (2.48)) is given by the commutator

$$[d_1, d_2] = d_1 d_2 - d_2 d_1. \tag{3.48}$$

Write

$$\mathcal{D}=\sum_{i=1}^{27}\mathbb{C}\partial_{x_i}. \tag{3.49}$$

Then $\mathcal{D}$ forms a $\mathcal{G}^{E_6}$-module with respect to the action

$$u(\partial)=[u|_{\mathcal{A}},\partial]\qquad\text{for }\ u\in\mathcal{G}^{E_6},\ \partial\in\mathcal{D}. \tag{3.50}$$

On the other hand, $\mathcal{G}_\pm$ (cf. (2.25)-(2.39)) form $\mathcal{G}^{E_6}$-modules with respect to the adjoint representation. According to (2.44) and (2.45), the linear map determined by $\eta_i\mapsto x_i$ for $i\in\overline{1,27}$ gives a $\mathcal{G}^{E_6}$-module isomorphism from $\mathcal{G}_-$ to V. Moreover,

$$(\xi_i|\eta_j)=-\delta_{i,j}\qquad\text{for }\ i,j\in\overline{1,27} \tag{3.51}$$

by (2.12). The fact $\partial_{x_i}(x_j)=\delta_{i,j}$ for $i,j\in\overline{1,27}$, (2.13) and (3.51) show that the linear map determined by $\xi_i\mapsto\partial_{x_i}$ for $i\in\overline{1,27}$ gives a $\mathcal{G}^{E_6}$-module isomorphism from $\mathcal{G}_+$ to $\mathcal{D}$. Hence we define the action of $\mathcal{G}_+$ on $\mathcal{A}$ by

$$\xi_i|_{\mathcal{A}}=\partial_{x_i}\qquad\text{for }\ i\in\overline{1,27}. \tag{3.52}$$

Recall the Witt Lie subalgebra of $\mathbb{A}$:

$$\mathcal{W}_{27}=\sum_{i=1}^{27}\mathcal{A}\partial_{x_i}. \tag{3.53}$$

Now we want to find the differential operators $P_1,P_2,...,P_{27}\in\mathcal{W}_{24}$ such that the following action matches the structure of $\mathcal{G}^{E_7}$:

$$\eta_i|_{\mathcal{A}}=P_i\qquad\text{for }\ i\in\overline{1,27}. \tag{3.54}$$

Comparing the weights in Table 1 and Table 2, we use (3.1) and (3.8)-(3.33) to assume

$$\begin{aligned}P_1 &= x_1D+c_1\zeta_1\partial_{x_{14}}+c_2\zeta_2\partial_{x_{17}}+c_3\zeta_3\partial_{x_{19}}+c_4\zeta_4\partial_{x_{21}}+c_5\zeta_5\partial_{x_{22}}\\ &\quad +c_6\zeta_6\partial_{x_{23}}+c_7\zeta_7\partial_{x_{24}}+c_9\zeta_9\partial_{x_{25}}+c_{11}\zeta_{11}\partial_{x_{26}}+c_{14}\zeta_{14}\partial_{x_{27}},\end{aligned} \tag{3.55}$$

where $c_i\in\mathbb{C}$. Imposing

$$\begin{aligned}[\partial_{x_1},P_1] &= [\xi_1,\eta_1]|_{\mathcal{A}}=-\alpha_7|_{\mathcal{A}}=2x_1\partial_{x_1}+\sum_{i=2}^{13}x_i\partial_{x_i}\\ &\quad +x_{15}\partial_{x_{15}}+x_{16}\partial_{x_{16}}+x_{18}\partial_{x_{18}}+x_{20}\partial_{x_{20}}\end{aligned} \tag{3.56}$$

by (2.87) and Table 1, we get

$$c_1=c_2=c_3=-c_4=c_5=c_6=-c_7=c_9=c_{11}=c_{14}=-1 \tag{3.57}$$

by (3.1), (3.8)-(3.13), (3.15), (3.17) and (3.20). Thus

$$\begin{aligned}P_1 &= x_1D-\zeta_1\partial_{x_{14}}-\zeta_2\partial_{x_{17}}-\zeta_3\partial_{x_{19}}+\zeta_4\partial_{x_{21}}-\zeta_5\partial_{x_{22}}\\ &\quad -\zeta_6\partial_{x_{23}}+\zeta_7\partial_{x_{24}}-\zeta_9\partial_{x_{25}}-\zeta_{11}\partial_{x_{26}}-\zeta_{14}\partial_{x_{27}}.\end{aligned} \tag{3.58}$$

According to (3.1), (3.8)-(3.13), (3.15), (3.17) and (3.20), we find

$$[\partial_{x_2},P_1]=x_1\partial_{x_2}-x_{11}\partial_{x_{14}}-x_{13}\partial_{x_{17}}-x_{16}\partial_{x_{19}}-x_{18}\partial_{x_{21}}-x_{20}\partial_{x_{23}}, \tag{3.59}$$

$$[\partial_{x_3},P_1]=x_1\partial_{x_3}-x_9\partial_{x_{14}}-x_{12}\partial_{x_{17}}-x_{15}\partial_{x_{19}}+x_{18}\partial_{x_{22}}+x_{20}\partial_{x_{24}}, \tag{3.60}$$

$$[\partial_{x_4},P_1]=x_1\partial_{x_4}-x_7\partial_{x_{14}}-x_{10}\partial_{x_{17}}+x_{15}\partial_{x_{21}}+x_{16}\partial_{x_{22}}-x_{20}\partial_{x_{25}}, \tag{3.61}$$

$$[\partial_{x_5},P_1]=x_1\partial_{x_5}+x_6\partial_{x_{14}}-x_{10}\partial_{x_{19}}-x_{12}\partial_{x_{21}}-x_{13}\partial_{x_{22}}+x_{20}\partial_{x_{25}}, \tag{3.62}$$

$$[\partial_{x_6},P_1]=x_1\partial_{x_6}+x_5\partial_{x_{14}}+x_8\partial_{x_{17}}-x_{15}\partial_{x_{23}}-x_{16}\partial_{x_{24}}-x_{18}\partial_{x_{25}}, \tag{3.63}$$

$$[\partial_{x_7},P_1]=x_1\partial_{x_7}-x_4\partial_{x_{14}}+x_8\partial_{x_{19}}+x_{12}\partial_{x_{23}}+x_{13}\partial_{x_{24}}+x_{18}\partial_{x_{26}}, \tag{3.64}$$

$$[\partial_{x_8},P_1]=x_1\partial_{x_8}+x_6\partial_{x_{17}}+x_7\partial_{x_{19}}+x_9\partial_{x_{21}}+x_{11}\partial_{x_{22}}+x_{20}\partial_{x_{27}}, \tag{3.65}$$

$$[\partial_{x_9},P_1]=x_1\partial_{x_9}-x_3\partial_{x_{14}}+x_8\partial_{x_{21}}-x_{10}\partial_{x_{23}}+x_{13}\partial_{x_{25}}-x_{16}\partial_{x_{26}}, \tag{3.66}$$

$$[\partial_{x_{10}},P_1]=x_1\partial_{x_{10}}-x_4\partial_{x_{17}}-x_5\partial_{x_{19}}-x_9\partial_{x_{23}}-x_{11}\partial_{x_{24}}+x_{18}\partial_{x_{27}}, \tag{3.67}$$

$$[\partial_{x_{11}},P_1]=x_1\partial_{x_{11}}-x_2\partial_{x_{14}}+x_8\partial_{x_{22}}-x_{10}\partial_{x_{24}}-x_{12}\partial_{x_{25}}+x_{15}\partial_{x_{26}}, \tag{3.68}$$

$$[\partial_{x_{12}},P_1]=x_1\partial_{x_{12}}-x_3\partial_{x_{17}}-x_5\partial_{x_{21}}+x_7\partial_{x_{23}}-x_{11}\partial_{x_{25}}-x_{16}\partial_{x_{27}}, \tag{3.69}$$

$$[\partial_{x_{13}},P_1]=x_1\partial_{x_{13}}-x_2\partial_{x_{17}}-x_5\partial_{x_{22}}+x_7\partial_{x_{24}}+x_9\partial_{x_{25}}+x_{15}\partial_{x_{27}}, \tag{3.70}$$

$$[\partial_{x_{15}},P_1]=x_1\partial_{x_{15}}-x_3\partial_{x_{19}}+x_4\partial_{x_{21}}-x_6\partial_{x_{23}}+x_{11}\partial_{x_{26}}+x_{13}\partial_{x_{27}}, \tag{3.71}$$

$$[\partial_{x_{16}},P_1]=x_1\partial_{x_{16}}-x_2\partial_{x_{19}}+x_4\partial_{x_{22}}-x_6\partial_{x_{24}}-x_9\partial_{x_{26}}-x_{12}\partial_{x_{27}}, \tag{3.72}$$

$$[\partial_{x_{18}},P_1]=x_1\partial_{x_{18}}-x_2\partial_{x_{21}}+x_3\partial_{x_{22}}-x_6\partial_{x_{25}}+x_7\partial_{x_{26}}+x_{10}\partial_{x_{27}}, \tag{3.73}$$

$$[\partial_{x_{20}},P_1]=x_1\partial_{x_{20}}-x_2\partial_{x_{23}}+x_3\partial_{x_{24}}-x_4\partial_{x_{25}}+x_5\partial_{x_{26}}+x_8\partial_{x_{27}}, \tag{3.74}$$

$$[x_r,P_1]=0\qquad\text{for }\ r=14,17,19,21,22,23,24,25,26,27. \tag{3.75}$$

On the other hand, if $[\xi_i,\eta_1]\neq 0$ for $i\geq 2$, then it is equal to the vector obtained by deleting the last coordinate 1 by (2.27)-(2.33). For instance, $\xi_{18}=E_{(1,1,2,3,2,1,1)}$ and $[\xi_{18},\eta_1]=E_{(1,1,2,3,2,1)}$. By (3.58)-(3.73) and correspondingly (2.55), (2.60), (2.65), (2.70), (2.69), (2.74), (2.72), (2.77), (2.76), (2.80), (2.79), (2.82), (2.81), (2.83)-(2.85), we have:

$$[\partial_{x_i},P_1]=[\xi_i,\eta_1]|_{\mathcal{A}}\qquad\text{for }\ i\in\overline{1,27}. \tag{3.76}$$

Expressions (2.50)-(2.55), (3.35)-(3.40) and (3.48) imply that P_1 is a $\mathcal{G}^{E_6}$-singular vector in $\mathcal{W}_{27}$ with weight λ_6. Since $-[E_{-\alpha_6},x_1]=x_2$, we set

$$\begin{aligned}P_2 &= -[E_{-\alpha_6},P_1]=x_2D-\zeta_1\partial_{x_{11}}-\zeta_2\partial_{x_{13}}-\zeta_3\partial_{x_{16}}+\zeta_4\partial_{x_{18}}\\ &\quad -\zeta_6\partial_{x_{20}}-\zeta_8\partial_{x_{22}}+\zeta_{10}\partial_{x_{24}}-\zeta_{12}\partial_{x_{25}}-\zeta_{15}\partial_{x_{26}}+\zeta_{17}\partial_{x_{27}}\end{aligned} \tag{3.77}$$

by (3.7), (3.46) and (3.58). As $-[E_{-\alpha_5},x_2]=x_3$, we take

$$\begin{aligned}P_3 &= -[E_{-\alpha_5},P_2]=x_3D-\zeta_1\partial_{x_9}-\zeta_2\partial_{x_{12}}-\zeta_3\partial_{x_{15}}+\zeta_5\partial_{x_{18}}\\ &\quad -\zeta_7\partial_{x_{20}}-\zeta_8\partial_{x_{21}}+\zeta_{10}\partial_{x_{23}}-\zeta_{13}\partial_{x_{25}}-\zeta_{16}\partial_{x_{26}}+\zeta_{19}\partial_{x_{27}}\end{aligned} \tag{3.78}$$

by (3.6), (3.45) and (3.77). Thanks to $-[E_{-\alpha_4},x_3]=x_4$, we have

$$\begin{aligned}P_4 &= -[E_{-\alpha_4},P_3]=x_4D-\zeta_1\partial_{x_7}-\zeta_2\partial_{x_{10}}-\zeta_4\partial_{x_{15}}+\zeta_5\partial_{x_{16}}\\ &\quad -\zeta_8\partial_{x_{19}}-\zeta_9\partial_{x_{20}}+\zeta_{12}\partial_{x_{23}}-\zeta_{13}\partial_{x_{24}}-\zeta_{18}\partial_{x_{26}}+\zeta_{21}\partial_{x_{27}}\end{aligned} \tag{3.79}$$

by (3.5), (3.44) and (3.78). Due to $-[E_{-\alpha_3},x_4]=x_5$, we write

$$\begin{aligned}P_5 &= -[E_{-\alpha_3},P_4]=x_5D+\zeta_1\partial_{x_6}-\zeta_3\partial_{x_{10}}+\zeta_4\partial_{x_{12}}-\zeta_5\partial_{x_{13}}\\ &\quad +\zeta_8\partial_{x_{17}}+\zeta_{11}\partial_{x_{20}}-\zeta_{15}\partial_{x_{23}}+\zeta_{16}\partial_{x_{24}}-\zeta_{18}\partial_{x_{25}}-\zeta_{22}\partial_{x_{27}}\end{aligned} \tag{3.80}$$

by (3.4), (3.43) and (3.79). Since $-[E_{-\alpha_2},x_4]=x_6$, we denote

$$\begin{aligned}P_6 &= -[E_{-\alpha_2},P_4]=x_6D+\zeta_1\partial_{x_5}+\zeta_2\partial_{x_8}-\zeta_6\partial_{x_{15}}+\zeta_7\partial_{x_{16}}\\ &\quad -\zeta_9\partial_{x_{18}}-\zeta_{10}\partial_{x_{19}}+\zeta_{12}\partial_{x_{21}}-\zeta_{13}\partial_{x_{22}}+\zeta_{20}\partial_{x_{26}}-\zeta_{23}\partial_{x_{27}}\end{aligned}\tag{3.81}$$

by (3.3), (3.42) and (3.79). As $-[E_{-\alpha_3},x_6]=x_7$, we denote

$$\begin{aligned}P_7 &= -[E_{-\alpha_3},P_6]=x_7D-\zeta_1\partial_{x_4}+\zeta_3\partial_{x_8}+\zeta_6\partial_{x_{12}}-\zeta_7\partial_{x_{13}}\\ &\quad +\zeta_{10}\partial_{x_{17}}+\zeta_{11}\partial_{x_{18}}-\zeta_{15}\partial_{x_{21}}+\zeta_{16}\partial_{x_{22}}+\zeta_{20}\partial_{x_{25}}+\zeta_{24}\partial_{x_{27}}\end{aligned}\tag{3.82}$$

by (3.4), (3.43) and (3.81). Thanks to $-[E_{-\alpha_1},x_5]=x_8$, we find

$$\begin{aligned}P_8 &= -[E_{-\alpha_2},P_5]=x_8D+\zeta_2\partial_{x_6}+\zeta_3\partial_{x_7}-\zeta_4\partial_{x_9}+\zeta_5\partial_{x_{11}}\\ &\quad -\zeta_8\partial_{x_{14}}+\zeta_{14}\partial_{x_{20}}+\zeta_{17}\partial_{x_{23}}-\zeta_{19}\partial_{x_{24}}+\zeta_{21}\partial_{x_{25}}+\zeta_{22}\partial_{x_{26}}\end{aligned}\tag{3.83}$$

by (3.2), (3.41) and (3.80).

Since $[E_{-\alpha_4},x_7]=x_9$, we take

$$\begin{aligned}P_9 &= [E_{-\alpha_4},P_7]=x_9D-\zeta_1\partial_{x_3}-\zeta_4\partial_{x_8}-\zeta_6\partial_{x_{10}}+\zeta_9\partial_{x_{13}}\\ &\quad -\zeta_{12}\partial_{x_{17}}-\zeta_{11}\partial_{x_{16}}+\zeta_{15}\partial_{x_{19}}-\zeta_{18}\partial_{x_{22}}-\zeta_{20}\partial_{x_{24}}+\zeta_{25}\partial_{x_{27}}\end{aligned}\tag{3.84}$$

by (3.5), (3.44) and (3.82). As $[E_{-\alpha_1},x_7]=-x_{10}$, we let

$$\begin{aligned}P_{10} &= -[E_{-\alpha_1},P_7]=x_{10}D-\zeta_2\partial_{x_4}-\zeta_3\partial_{x_5}-\zeta_6\partial_{x_9}+\zeta_7\partial_{x_{11}}\\ &\quad -\zeta_{10}\partial_{x_{14}}+\zeta_{14}\partial_{x_{18}}+\zeta_{17}\partial_{x_{21}}-\zeta_{19}\partial_{x_{22}}-\zeta_{23}\partial_{x_{25}}-\zeta_{24}\partial_{x_{26}}\end{aligned}\tag{3.85}$$

by (3.2), (3.41) and (3.82). Due to $[E_{-\alpha_5},x_9]=x_{11}$, we take

$$\begin{aligned}P_{11} &= [E_{-\alpha_4},P_7]=x_{11}D-\zeta_1\partial_{x_2}+\zeta_5\partial_{x_8}+\zeta_7\partial_{x_{10}}-\zeta_9\partial_{x_{12}}\\ &\quad +\zeta_{11}\partial_{x_{15}}+\zeta_{13}\partial_{x_{17}}-\zeta_{16}\partial_{x_{19}}+\zeta_{18}\partial_{x_{21}}+\zeta_{20}\partial_{x_{23}}+\zeta_{26}\partial_{x_{27}}\end{aligned}\tag{3.86}$$

by (3.6), (3.45) and (3.84). Thanks to $-[E_{-\alpha_1},x_9]=x_{12}$, we find

$$\begin{aligned}P_{12} &= -[E_{-\alpha_1},P_9]=x_{12}D-\zeta_2\partial_{x_3}+\zeta_4\partial_{x_5}+\zeta_6\partial_{x_7}-\zeta_9\partial_{x_{11}}\\ &\quad +\zeta_{12}\partial_{x_{14}}-\zeta_{14}\partial_{x_{16}}-\zeta_{17}\partial_{x_{19}}+\zeta_{21}\partial_{x_{22}}+\zeta_{23}\partial_{x_{24}}-\zeta_{25}\partial_{x_{26}}\end{aligned}\tag{3.87}$$

by (3.2), (3.41) and (3.84). Due to $[E_{-\alpha_5},x_{12}]=x_{13}$, we write

$$\begin{aligned}P_{13} &= [E_{-\alpha_5},P_{12}]=x_{13}D-\zeta_2\partial_{x_2}-\zeta_5\partial_{x_5}-\zeta_7\partial_{x_7}+\zeta_9\partial_{x_9}\\ &\quad -\zeta_{13}\partial_{x_{14}}+\zeta_{14}\partial_{x_{15}}+\zeta_{19}\partial_{x_{19}}-\zeta_{21}\partial_{x_{21}}-\zeta_{23}\partial_{x_{23}}-\zeta_{26}\partial_{x_{26}}\end{aligned}\tag{3.88}$$

by (3.6), (3.45) and (3.87). Equation $[E_{-\alpha_6},x_{11}]=x_{14}$, we let

$$\begin{aligned}P_{14} &= [E_{-\alpha_6},P_{11}]=x_{14}D-\zeta_1\partial_{x_1}-\zeta_8\partial_{x_8}-\zeta_{10}\partial_{x_{10}}+\zeta_{12}\partial_{x_{12}}\\ &\quad -\zeta_{13}\partial_{x_{13}}-\zeta_{15}\partial_{x_{15}}+\zeta_{16}\partial_{x_{16}}-\zeta_{18}\partial_{x_{18}}-\zeta_{20}\partial_{x_{20}}+\zeta_{27}\partial_{x_{27}}\end{aligned}\tag{3.89}$$

by (3.7), (3.46) and (3.86).

As $-[E_{-\alpha_3},x_{12}]=x_{15}$, we get

$$\begin{aligned}P_{15} &= -[E_{-\alpha_3},P_{12}]=x_{15}D-\zeta_3\partial_{x_3}-\zeta_4\partial_{x_4}-\zeta_6\partial_{x_6}+\zeta_{11}\partial_{x_{11}}\\ &\quad +\zeta_{14}\partial_{x_{13}}-\zeta_{15}\partial_{x_{14}}+\zeta_{17}\partial_{x_{17}}-\zeta_{22}\partial_{x_{22}}-\zeta_{24}\partial_{x_{24}}-\zeta_{25}\partial_{x_{25}}\end{aligned}\tag{3.90}$$

by (3.4), (3.43) and (3.87). Thanks to $[E_{-\alpha_5}, x_{15}] = x_{16}$, we let

$$\begin{aligned} P_{16} &= [E_{-\alpha_5}, P_{15}] = x_{16}D - \zeta_3\partial_{x_2} + \zeta_5\partial_{x_4} + \zeta_7\partial_{x_6} - \zeta_{11}\partial_{x_9} \\ &\quad -\zeta_{14}\partial_{x_{12}} + \zeta_{16}\partial_{x_{14}} - \zeta_{19}\partial_{x_{17}} + \zeta_{22}\partial_{x_{21}} + \zeta_{24}\partial_{x_{23}} - \zeta_{26}\partial_{x_{25}} \end{aligned} \tag{3.91}$$

by (3.6), (3.45) and (3.90). Due to $-[E_{-\alpha_1}, x_{14}] = x_{17}$, we get

$$\begin{aligned} P_{17} &= = -[E_{-\alpha_1}, P_{14}] = x_{17}D - \zeta_2\partial_{x_1} + \zeta_8\partial_{x_5} + \zeta_{10}\partial_{x_7} - \zeta_{12}\partial_{x_9} \\ &\quad +\zeta_{13}\partial_{x_{11}} + \zeta_{17}\partial_{x_{15}} - \zeta_{19}\partial_{x_{16}} + \zeta_{21}\partial_{x_{18}} + \zeta_{23}\partial_{x_{20}} - \zeta_{27}\partial_{x_{26}} \end{aligned} \tag{3.92}$$

by (3.2), (3.41) and (3.89). Since $[E_{-\alpha_4}, x_{16}] = x_{18}$, we find

$$\begin{aligned} P_{18} &= [E_{-\alpha_4}, P_{16}] = x_{18}D + \zeta_4\partial_{x_2} + \zeta_5\partial_{x_3} - \zeta_9\partial_{x_6} + \zeta_{11}\partial_{x_7} \\ &\quad +\zeta_{14}\partial_{x_{10}} - \zeta_{18}\partial_{x_{14}} + \zeta_{21}\partial_{x_{17}} - \zeta_{22}\partial_{x_{19}} + \zeta_{25}\partial_{x_{23}} + \zeta_{26}\partial_{x_{24}} \end{aligned} \tag{3.93}$$

by (3.5), (3.44) and (3.91).

Thanks to $[E_{-\alpha_6}, x_{16}] = x_{19}$, we calculate

$$\begin{aligned} P_{19} &= [E_{-\alpha_6}, P_{16}] = x_{19}D - \zeta_3\partial_{x_1} - \zeta_8\partial_{x_4} - \zeta_{10}\partial_{x_6} + \zeta_{15}\partial_{x_9} \\ &\quad -\zeta_{17}\partial_{x_{12}} - \zeta_{16}\partial_{x_{11}} + \zeta_{19}\partial_{x_{13}} - \zeta_{22}\partial_{x_{18}} - \zeta_{24}\partial_{x_{20}} - \zeta_{27}\partial_{x_{25}} \end{aligned} \tag{3.94}$$

by (3.7), (3.46) and (3.91). As $[E_{-\alpha_2}, x_{18}] = x_{20}$, we have

$$\begin{aligned} P_{20} &= [E_{-\alpha_2}, P_{18}] = x_{20}D - \zeta_6\partial_{x_2} - \zeta_7\partial_{x_3} - \zeta_9\partial_{x_4} + \zeta_{11}\partial_{x_5} \\ &\quad +\zeta_{14}\partial_{x_8} - \zeta_{20}\partial_{x_{14}} + \zeta_{23}\partial_{x_{17}} - \zeta_{24}\partial_{x_{19}} - \zeta_{25}\partial_{x_{21}} - \zeta_{26}\partial_{x_{22}} \end{aligned} \tag{3.95}$$

by (3.3), (3.42) and (3.93). Since $[E_{-\alpha_4}, x_{19}] = x_{21}$, we let

$$\begin{aligned} P_{21} &= [E_{-\alpha_4}, P_{19}] = x_{21}D + \zeta_4\partial_{x_1} - \zeta_8\partial_{x_3} + \zeta_{12}\partial_{x_6} - \zeta_{15}\partial_{x_7} \\ &\quad +\zeta_{17}\partial_{x_{10}} + \zeta_{18}\partial_{x_{11}} - \zeta_{21}\partial_{x_{13}} + \zeta_{22}\partial_{x_{16}} - \zeta_{25}\partial_{x_{20}} + \zeta_{27}\partial_{x_{24}} \end{aligned} \tag{3.96}$$

by (3.5), (3.44) and (3.94). Equation $[E_{-\alpha_5}, x_{21}] = x_{22}$, we write

$$\begin{aligned} P_{22} &= [E_{-\alpha_5}, P_{21}] = x_{22}D - \zeta_5\partial_{x_1} - \zeta_8\partial_{x_2} - \zeta_{13}\partial_{x_6} + \zeta_{16}\partial_{x_7} \\ &\quad -\zeta_{19}\partial_{x_{10}} - \zeta_{18}\partial_{x_9} + \zeta_{21}\partial_{x_{12}} - \zeta_{22}\partial_{x_{15}} - \zeta_{26}\partial_{x_{20}} - \zeta_{27}\partial_{x_{23}} \end{aligned} \tag{3.97}$$

by (3.6), (3.45) and (3.96).

Note that the equation $[E_{-\alpha_2}, x_{21}] = x_{23}$ gives rise to

$$\begin{aligned} P_{23} &= [E_{-\alpha_2}, P_{21}] = x_{23}D - \zeta_6\partial_{x_1} + \zeta_{10}\partial_{x_3} + \zeta_{12}\partial_{x_4} - \zeta_{15}\partial_{x_5} \\ &\quad +\zeta_{17}\partial_{x_8} + \zeta_{20}\partial_{x_{11}} - \zeta_{23}\partial_{x_{13}} + \zeta_{24}\partial_{x_{16}} + \zeta_{25}\partial_{x_{18}} - \zeta_{27}\partial_{x_{22}} \end{aligned} \tag{3.98}$$

by (3.3), (3.42) and (3.96). Since $[E_{-\alpha_2}, x_{22}] = x_{24}$, we write

$$\begin{aligned} P_{24} &= [E_{-\alpha_2}, P_{22}] = x_{24}D + \zeta_7\partial_{x_1} + \zeta_{10}\partial_{x_2} - \zeta_{13}\partial_{x_4} + \zeta_{16}\partial_{x_5} \\ &\quad -\zeta_{19}\partial_{x_8} - \zeta_{20}\partial_{x_9} + \zeta_{23}\partial_{x_{12}} - \zeta_{24}\partial_{x_{15}} + \zeta_{26}\partial_{x_{18}} + \zeta_{27}\partial_{x_{21}} \end{aligned} \tag{3.99}$$

by (3.3), (3.42) and (3.97). As $[E_{-\alpha_4}, x_{24}] = x_{25}$, we write

$$\begin{aligned} P_{25} &= [E_{-\alpha_4}, P_{24}] = x_{25}D - \zeta_9\partial_{x_1} - \zeta_{12}\partial_{x_2} - \zeta_{13}\partial_{x_3} - \zeta_{18}\partial_{x_5} \\ &\quad +\zeta_{21}\partial_{x_8} + \zeta_{20}\partial_{x_7} - \zeta_{23}\partial_{x_{10}} - \zeta_{25}\partial_{x_{15}} - \zeta_{26}\partial_{x_{16}} - \zeta_{27}\partial_{x_{19}} \end{aligned} \tag{3.100}$$

by (3.5), (3.44) and (3.99). Thanks to $[E_{-\alpha_3}, x_{25}] = x_{26}$, we write

$$\begin{aligned} P_{26} &= [E_{-\alpha_3}, P_{25}] = x_{26}D - \zeta_{11}\partial_{x_1} - \zeta_{15}\partial_{x_2} - \zeta_{16}\partial_{x_3} - \zeta_{18}\partial_{x_4} \\ &\quad +\zeta_{22}\partial_{x_8} + \zeta_{20}\partial_{x_6} - \zeta_{24}\partial_{x_{10}} - \zeta_{25}\partial_{x_{12}} - \zeta_{26}\partial_{x_{13}} - \zeta_{27}\partial_{x_{17}} \end{aligned} \tag{3.101}$$

by (3.4), (3.43) and (3.100). Due to $-[E_{-\alpha_1}, x_{26}] = x_{27}$, we write

$$\begin{aligned} P_{27} &= -[E_{-\alpha_1}, P_{26}] = x_{27}D - \zeta_{14}\partial_{x_1} + \zeta_{17}\partial_{x_2} + \zeta_{19}\partial_{x_3} + \zeta_{21}\partial_{x_4} \\ &\quad -\zeta_{22}\partial_{x_5} - \zeta_{23}\partial_{x_6} + \zeta_{24}\partial_{x_7} + \zeta_{25}\partial_{x_9} + \zeta_{26}\partial_{x_{11}} + \zeta_{27}\partial_{x_{14}} \end{aligned} \tag{3.102}$$

by (3.2), (3.41) and (3.101).

Set

$$\mathcal{P} = \sum_{i=1}^{27} \mathbb{C}P_i, \qquad \mathcal{C}_0 = \mathcal{G}^{E_6}|_{\mathcal{A}} + \mathbb{C}D \tag{3.103}$$

(cf. (2.50)-(2.87) and (2.91)) and

$$\mathcal{C} = \mathcal{P} + \mathcal{C}_0 + \mathcal{D} \tag{3.104}$$

(cf. (3.49)). Then we have:

Theorem 3.1. *The space $\mathcal{C}$ forms a Lie subalgebra of the Witt algebra $\mathcal{W}_{27}$ (cf. (3.53)). Moreover, the linear map ϑ determined by*

$$\vartheta(\xi_i) = \partial_{x_i}, \ \ \vartheta(\eta_i) = P_i, \ \ \vartheta(u) = u|_{\mathcal{A}} \qquad \text{for } \ i \in \overline{1,27}, \ u \in \mathcal{G}^{E_6} \tag{3.105}$$

and

$$\vartheta(\alpha_7) = -2x_1\partial_{x_1} - \sum_{i=2}^{13} x_i\partial_{x_i} - x_{15}\partial_{x_{15}} - x_{16}\partial_{x_{16}} - x_{18}\partial_{x_{18}} - x_{20}\partial_{x_{20}} \tag{3.106}$$

(cf. (2.87) and Table 1) gives a Lie algebra isomorphism from $\mathcal{G}^{E_7}$ to $\mathcal{C}$.

Proof. Since $\mathcal{D} \cong \mathcal{G}_+$ as $\mathcal{G}^{E_6}$-modules, we have

$$\mathcal{G}_0 + \mathcal{G}_+ \overset{\vartheta}{\cong} \mathcal{C}_0 + \mathcal{D} \tag{3.107}$$

as Lie algebras. Denote by $U(\mathcal{G})$ the universal enveloping algebra of a Lie algebra $\mathcal{G}$. Note that

$$\mathcal{B}_- = \mathcal{G}_0 + \mathcal{G}_-, \qquad \mathcal{B}_+ = \mathcal{G}_0 + \mathcal{G}_+ \tag{3.108}$$

are also Lie subalgebras of $\mathcal{G}^{E_7}$ and

$$\mathcal{G}^{E_7} = \mathcal{B}_- \oplus \mathcal{G}_+ = \mathcal{G}_- \oplus \mathcal{B}_+. \tag{3.109}$$

We define a one-dimensional $\mathcal{B}_-$-module $\mathbb{C}u_0$ by

$$w(u_0) = 0 \qquad \text{for } \ w \in \mathcal{B}_- \bigcap \mathcal{G}^{E_6}, \ \ \widehat{\alpha}(u_0) = -27u_0 \tag{3.110}$$

(cf. (2.88)). Let

$$\Psi = U(\mathcal{G}^{E_7}) \otimes_{\mathcal{B}_-} \mathbb{C}u_0 \cong U(\mathcal{G}_+) \otimes_{\mathbb{C}} \mathbb{C}u_0 \tag{3.111}$$

be the induced $\mathcal{G}^{E_7}$-module.

Recall that $\mathbb{N}$ is the set of nonnegative integers. Let

$$\mathcal{A}' = \mathbb{C}[\partial_{x_1}, \partial_{x_2}, ..., \partial_{x_{27}}]. \tag{3.112}$$

We define an action of the associative algebra $\mathbb{A}$ (cf. (2.48)) on $\mathcal{A}'$ by

$$\partial_{x_i}(\prod_{j=1}^{27} \partial_{x_j}^{\beta_j}) = \partial_{x_i}^{\beta_i+1} \prod_{i\neq j\in\overline{1,27}} \prod_{j=1}^{27} \partial_{x_j}^{\beta_j} \tag{3.113}$$

and

$$x_i(\prod_{j=1}^{27} \partial_{x_j}^{\beta_j}) = -\beta_i \partial_{x_i}^{\beta_i-1} \prod_{i\neq j\in\overline{1,27}} \prod_{j=1}^{27} \partial_{x_j}^{\beta_j} \tag{3.114}$$

for $i \in \overline{1,27}$. Since

$$[-x_i, \partial_{x_j}] = [\partial_{x_i}, x_j] = \delta_{i,j} \qquad \text{for } \ i,j \in \overline{1,27}, \tag{3.115}$$

the above action gives an associative algebra representation of $\mathbb{A}$. Thus it also gives a Lie algebra representation of $\mathbb{A}$ (cf. (3.48)). It is straightforward to verify that

$$[d|_{\mathcal{A}'}, \partial|_{\mathcal{A}'}] = [d, \partial]|_{\mathcal{A}'} \qquad \text{for } \ d \in \mathcal{C}_0, \ \partial \in \mathcal{D}. \tag{3.116}$$

Define linear map $\varsigma : \Psi \to \mathcal{A}'$ by

$$\varsigma(\prod_{i=1}^{27} \xi_i^{\ell_i} \otimes u_0) = \prod_{i=1}^{27} \partial_{x_i}^{\ell_i} \qquad (\ell_1, ..., \ell_{27}) \in \mathbb{N}^{27}. \tag{3.117}$$

According to (2.50)-(2.87), (3.113) and (3.114),

$$D(1) = -27, \ \ d(1) = 0 \qquad \text{for } \ d \in o(10,\mathbb{C})|_{\mathcal{A}}. \tag{3.118}$$

Moreover, (3.110), (3.111), (3.113), (3.114) and (3.118) imply

$$\varsigma(\xi(v)) = \vartheta(\xi)\varsigma(v) \qquad \text{for } \ \xi \in \mathcal{G}_0, \ v \in \Psi. \tag{3.119}$$

Now (3.111) and (3.113) imply

$$\varsigma(w(u)) = \vartheta(w)(\varsigma(u)) \qquad \text{for } \ w \in \mathcal{B}_+, \ \ u \in \Psi. \tag{3.120}$$

Thus we have

$$\varsigma w|_{\Psi}\varsigma^{-1} = \vartheta(w)|_{\mathcal{A}'} \qquad \text{for } \ w \in \mathcal{B}_+. \tag{3.121}$$

On the other hand, the linear map

$$\psi(v) = \varsigma v|_{\Psi}\varsigma^{-1} \qquad \text{for } \ v \in \mathcal{G}^{E_7} \tag{3.122}$$

is a Lie algebra monomorphism from $\mathcal{G}^{E_7}$ to $\mathbb{A}|_{\mathcal{A}'}$. According to (3.76) and (3.115),

$$\psi(\eta_1) = P_1|_{\mathcal{A}'}. \tag{3.123}$$

By the constructions of $P_2, ..., P_{27}$ in (2.77)-(2.102), we have

$$\psi(\eta_i) = P_i|_{\mathcal{A}'} \qquad \text{for } \ i \in \overline{2,27}. \tag{3.124}$$

Therefore, we have

$$\psi(v)=\vartheta(v)|_{\mathcal{A}'}\qquad\text{for}\ \ v\in\mathcal{G}^{E_7}. \tag{3.125}$$

In particular, $\mathcal{C}|_{\mathcal{A}'}=\vartheta(\mathcal{G}^{E_7})|_{\mathcal{A}'}=\psi(\mathcal{G}^{E_7})$ forms a Lie algebra. Since the linear map $d\mapsto d|_{\mathcal{A}'}$ for $d\in\mathcal{C}$ is injective, we have that $\mathcal{C}$ forms a Lie subalgebra of $\mathbb{A}$ and ϑ is a Lie algebra isomorphism. $\Box$

By the above theorem, a Lie group of type E_7 is generated by the linear transformations $\{e^{bu|_{\mathcal{A}}}\mid b\in\mathbb{R},\ u\in\mathcal{G}^{E_6}\}$ associated with (2.50)-(2.87), the real translations and dilations in $\sum_{i=1}^{27}\mathbb{R}x_i$, and the fractional transformations $\{e^{bP_i}\mid b\in\mathbb{R},\ i\in\overline{1,27}\}$ such as

$$e^{bP_1}(x_i)=\frac{x_i}{1-bx_1},\ \ i\in\{\overline{1,13},15,16,18,20\}, \tag{3.126}$$

$$e^{bP_1}(x_{14})=x_{14}-\frac{b(x_2x_{11}+x_3x_9+x_4x_7-x_5x_6)}{1-bx_1}, \tag{3.127}$$

$$e^{bP_1}(x_{17})=x_{17}-\frac{b(x_2x_{13}+x_3x_{12}+x_4x_{10}-x_6x_8)}{1-bx_1}, \tag{3.128}$$

$$e^{bP_1}(x_{19})=x_{19}-\frac{b(x_2x_{16}+x_3x_{15}+x_5x_{10}-x_7x_8)}{1-bx_1}, \tag{3.129}$$

$$e^{bP_1}(x_{21})=x_{21}-\frac{b(x_2x_{18}-x_4x_{15}+x_5x_{12}-x_8x_9)}{1-bx_1}, \tag{3.130}$$

$$e^{bP_1}(x_{22})=x_{22}+\frac{b(x_3x_{18}+x_4x_{16}-x_5x_{13}+x_8x_{11})}{1-bx_1}, \tag{3.131}$$

$$e^{bP_1}(x_{23})=x_{23}-\frac{b(x_2x_{20}+x_6x_{15}-x_7x_{12}+x_9x_{10})}{1-bx_1}, \tag{3.132}$$

$$e^{bP_1}(x_{24})=x_{24}+\frac{b(x_3x_{20}-x_6x_{16}+x_7x_{13}-x_{10}x_{11})}{1-bx_1}, \tag{3.133}$$

$$e^{bP_1}(x_{25})=x_{25}-\frac{b(x_4x_{20}+x_6x_{18}-x_9x_{13}+x_{11}x_{12})}{1-bx_1}, \tag{3.134}$$

$$e^{bP_1}(x_{26})=x_{26}+\frac{b(x_5x_{20}+x_7x_{18}-x_9x_{16}+x_{11}x_{15})}{1-bx_1}, \tag{3.135}$$

$$e^{bP_1}(x_{27})=x_{27}+\frac{b(x_8x_{20}+x_{10}x_{18}-x_{12}x_{16}+x_{13}x_{15})}{1-bx_1} \tag{3.136}$$

by (3.1)-(3.7), (3.9), (3.11), (3.14) and (3.58).

Later, we need the following Dickson's $\mathcal{G}^{E_6}$-invariant:

$$\begin{aligned}\chi\ =\ & x_1(x_{17}x_{26}+x_{19}x_{25}-x_{21}x_{24}+x_{22}x_{23})-x_{14}(x_1x_{27}-x_8x_{20}-x_{10}x_{18}+x_{12}x_{16}\\ &-x_{13}x_{15})+x_2(x_{13}x_{26}+x_{16}x_{25}-x_{18}x_{24}+x_{20}x_{22})-x_{11}(x_2x_{27}+x_8x_{23}+x_{10}x_{21}\\ &-x_{12}x_{19}+x_{15}x_{17})+x_3(x_{12}x_{26}+x_{15}x_{25}-x_{18}x_{23}+x_{20}x_{21})-x_9(x_3x_{27}-x_8x_{24}\\ &-x_{10}x_{22}+x_{13}x_{19}-x_{16}x_{17})+x_4(x_{10}x_{26}+x_{15}x_{24}-x_{16}x_{23}+x_{19}x_{20})-x_7(x_4x_{27}\\ &+x_8x_{25}+x_{12}x_{22}-x_{13}x_{21}+x_{17}x_{18})+x_5(x_{10}x_{25}-x_{12}x_{24}+x_{13}x_{23}-x_{17}x_{20})\\ &+x_6(x_5x_{27}-x_8x_{26}+x_{15}x_{22}-x_{16}x_{21}+x_{18}x_{19}).\end{aligned} \tag{3.137}$$

In [X2], we used partial differential equations to prove that any $\mathcal{G}^{E_6}$-singular vector in $\mathcal{A}$ is a polynomial in $x_1,\ \zeta_1,\ \chi$. Moreover,

$$\sum_{14\neq i\in\overline{1,26}} x_i\zeta_{28-i}-x_{14}\zeta_{14}-x_{27}\zeta_1=-3\chi \tag{3.138}$$

by Table 1, Table 2, (2.50)-(2.55), (3.1), (3.8)-(3.33) and (3.35)-(3.40).

4 Functor from E_6-Mod to E_7-Mod

In this section, we construct a new functor from E_6-**Mod** to E_7-**Mod**.

Note that

$$\mathcal{G}^{E_6}_{\mathcal{A}}=(\bigoplus_{i=1}^{6}\mathcal{A}\alpha_i)\oplus\bigoplus_{\alpha\in\Phi_{E_6}}\mathcal{A}E_\alpha \tag{4.1}$$

forms a Lie algebra with the Lie bracket:

$$[fu_1,gu_2]=fg[u_1,u_2]\qquad\text{for }\ f,g\in\mathcal{A},\ u_1,u_2\in\mathcal{G}^{E_6}. \tag{4.2}$$

Moreover, we define the Lie algebra

$$\mathcal{K}=\mathcal{G}^{E_6}_{\mathcal{A}}\oplus\mathcal{A}\kappa \tag{4.3}$$

with the Lie bracket:

$$[\xi_1+f\kappa,\xi_2+g\kappa]=[\xi_1,\xi_2]\qquad\text{for }\ \xi_1,\xi_2\in\mathcal{G}^{E_6}_{\mathcal{A}},\ f,g\in\mathcal{A}. \tag{4.4}$$

Similarly, $gl(27,\mathcal{A})$ becomes a Lie algebra with the Lie bracket as that in (4.2). Recall the Witt algebra $\mathcal{W}_{27}=\sum_{i=1}^{27}\mathcal{A}\partial_{x_i}$, and Shen [Sg1-3] found a monomorphism $\Im$ from the Lie algebra $\mathcal{W}_{27}$ to the Lie algebra of semi-product $\mathcal{W}_{27}+gl(27,\mathcal{A})$ defined by

$$\Im(\sum_{i=1}^{27}f_i\partial_{x_i})=\sum_{i=1}^{27}f_i\partial_{x_i}+\Im_1(\sum_{i=1}^{27}f_i\partial_{x_i}),\ \ \Im_1(\sum_{i=1}^{27}f_i\partial_{x_i})=\sum_{i,j=1}^{27}\partial_{x_i}(f_j)E_{i,j}. \tag{4.5}$$

According to our construction of P_1-P_{27} in (3.58) and (3.77)-(3.102),

$$\Im_1(P_i)=\sum_{r=1}^{27}x_r\Im_1([\xi_r,\eta_i]|_{\mathcal{A}})\qquad\text{for }\ i\in\overline{1,27}. \tag{4.6}$$

On the other hand,

$$\widehat{\mathcal{K}}=\mathcal{W}_{27}\oplus\mathcal{K} \tag{4.7}$$

becomes a Lie algebra with the Lie bracket

$$\begin{aligned}&[d_1+f_1u_1+f_2\kappa,d_2+g_1u_2+g_2\kappa]\\ =\ &[d_1,d_2]+f_1g_1[u_1,u_2]+d_1(g_2)u_2-d_2(g_1)u_1+(d_1(g_2)-d_2(g_1))\kappa\end{aligned} \tag{4.8}$$

for $f_1,f_2,g_1,g_2\in\mathcal{A},\ u_1,u_2\in\mathcal{G}^{E_6}$ and $d_1,d_2\in\mathcal{W}_{27}$. Note

$$\mathcal{G}_0=\mathcal{G}^{E_6}\oplus\mathbb{C}\widehat{\alpha} \tag{4.9}$$

by (2.41) and (2.88). So there exists a Lie algebra monomorphism $\varrho : \mathcal{G}_0 \to \mathcal{K}$ determined by

$$\varrho(\widehat{\alpha}) = 3\kappa, \quad \varrho(u) = u \qquad \text{for } u \in \mathcal{G}^{E_6}. \tag{4.10}$$

Since $\Im$ is a Lie algebra monomorphism, our construction of P_1-P_{27} in (3.48) and (3.58)-(3.102) show that we have a Lie algebra monomorphism $\iota : \mathcal{G}^{E_7} \to \widehat{\mathcal{K}}$ given by

$$\iota(u) = u|_{\mathcal{A}} + \varrho(u) \qquad \text{for } u \in \mathcal{G}_0, \tag{4.11}$$

$$\iota(\xi_i) = \partial_{x_i}, \quad \iota(\eta_i) = P_i + \sum_{r=1}^{27} x_r \varrho([\xi_r, \eta_i]) \qquad \text{for } i \in \overline{1,27}. \tag{4.12}$$

According to (2.4), (2.11) and (2.34)-(2.40),

$$\begin{aligned} \iota(\eta_1) \;=\;& P_1 + x_2 E_{\alpha_6} + x_3 E_{(0,0,0,0,1,1)} + x_4 E_{(0,0,0,1,1,1)} + x_5 E_{(0,0,1,1,1,1)} + x_6 E_{(0,1,0,1,1,1)} \\ & - x_1 \varrho(\alpha_7) + x_7 E_{(0,1,1,1,1,1)} + x_8 E_{(1,0,1,1,1,1)} + x_9 E_{(0,1,1,2,1,1)} + x_{10} E_{(1,1,1,1,1,1)} \\ & + x_{11} E_{(0,1,1,2,2,1)} + x_{12} E_{(1,1,1,2,1,1)} + x_{13} E_{(1,1,1,2,2,1)} + x_{15} E_{(1,1,2,2,1,1)} \\ & + x_{16} E_{(1,1,2,2,2,1)} + x_{18} E_{(1,1,2,3,2,1)} + x_{20} E_{(1,2,2,3,2,1)}. \end{aligned} \tag{4.13}$$

Moreover, (3.7) yields

$$\begin{aligned} \iota(\eta_2) \;=\;& -\iota([E_{-\alpha_6}, \eta_1]) = -[\iota(E_{-\alpha_6}), \iota(\eta_1)] = -[E_{-\alpha_6}|_{\mathcal{A}} + E_{-\alpha_6}, \iota(\eta_1)] \\ =\;& P_2 - x_1 E_{-\alpha_6} + x_3 E_{\alpha_5} + x_4 E_{(0,0,0,1,1)} + x_5 E_{(0,0,1,1,1)} + x_6 E_{(0,1,0,1,1)} \\ & - x_2 \varrho(\alpha_6 + \alpha_7) + x_7 E_{(0,1,1,1,1)} + x_8 E_{(1,0,1,1,1)} + x_9 E_{(0,1,1,2,1)} + x_{10} E_{(1,1,1,1,1)} \\ & + x_{12} E_{(1,1,1,2,1)} - x_{14} E_{(0,1,1,2,2,1)} + x_{15} E_{(1,1,2,2,1)} - x_{17} E_{(1,1,1,2,2,1)} \\ & - x_{19} E_{(1,1,2,2,2,1)} - x_{21} E_{(1,1,2,3,2,1)} - x_{23} E_{(1,2,2,3,2,1)}. \end{aligned} \tag{4.14}$$

Expression (3.6) gives

$$\begin{aligned} \iota(\eta_3) \;=\;& -\iota([E_{-\alpha_5}, \eta_2]) = -[\iota(E_{-\alpha_5}), \iota(\eta_2)] = -[E_{-\alpha_5}|_{\mathcal{A}} + E_{-\alpha_5}, \iota(\eta_2)] \\ =\;& P_3 - x_1 E'_{(0,0,0,0,1,1)} - x_2 E_{-\alpha_5} + x_4 E_{\alpha_4} + x_5 E_{(0,0,1,1)} + x_6 E_{(0,1,0,1)} \\ & - x_3 \varrho(\alpha_5 + \alpha_6 + \alpha_7) + x_7 E_{(0,1,1,1)} + x_8 E_{(1,0,1,1)} + x_{10} E_{(1,1,1,1)} - x_{11} E_{(0,1,1,2,1)} \\ & - x_{13} E_{(1,1,1,2,1)} - x_{14} E_{(0,1,1,2,1,1)} - x_{16} E_{(1,1,2,2,1)} - x_{17} E_{(1,1,1,2,1,1)} \\ & - x_{19} E_{(1,1,2,2,1,1)} + x_{22} E_{(1,1,2,3,2,1)} + x_{24} E_{(1,2,2,3,2,1)}. \end{aligned} \tag{4.15}$$

Furthermore,

$$\begin{aligned} \iota(\eta_4) \;=\;& -\iota([E_{-\alpha_4}, \eta_3]) = -[\iota(E_{-\alpha_4}), \iota(\eta_3)] = -[E_{-\alpha_4}|_{\mathcal{A}} + E_{-\alpha_4}, \iota(\eta_3)] \\ =\;& P_4 - x_1 E'_{(0,0,0,1,1,1)} - x_2 E'_{(0,0,0,1,1)} - x_3 E_{-\alpha_4} + x_5 E_{\alpha_3} + x_6 E_{\alpha_2} \\ & - x_4 \varrho(\sum_{i=4}^{7} \alpha_i) - x_9 E_{(0,1,1,1)} + x_8 E_{(1,0,1)} - x_{12} E_{(1,1,1,1)} - x_{11} E_{(0,1,1,1,1)} \\ & - x_{13} E_{(1,1,1,1,1)} - x_{14} E_{(0,1,1,1,1,1)} - x_{17} E_{(1,1,1,1,1,1)} + x_{18} E_{(1,1,2,2,1)} \\ & + x_{21} E_{(1,1,2,2,1,1)} + x_{22} E_{(1,1,2,2,2,1)} - x_{25} E_{(1,2,2,3,2,1)} \end{aligned} \tag{4.16}$$

by (3.5),

$$\begin{aligned}
\iota(\eta_5) &= -\iota([E_{-\alpha_3}, \eta_4]) = -[\iota(E_{-\alpha_3}), \iota(\eta_4)] = -[E_{-\alpha_3}|_{\mathcal{A}} + E_{-\alpha_3}, \iota(\eta_4)] \\
&= P_5 - x_1 E'_{(0,0,1,1,1,1)} - x_2 E'_{(0,0,1,1,1)} - x_3 E'_{(0,0,1,1)} - x_4 E_{-\alpha_3} + x_7 E_{\alpha_2} \\
&\quad -x_5 \varrho(\sum_{i=3}^{7} \alpha_i) + x_8 E_{\alpha_1} + x_9 E_{(0,1,0,1)} - x_{15} E_{(1,1,1,1)} + x_{11} E_{(0,1,0,1,1)} \\
&\quad -x_{16} E_{(1,1,1,1,1)} + x_{14} E_{(0,1,0,1,1,1)} - x_{19} E_{(1,1,1,1,1,1)} - x_{18} E_{(1,1,1,2,1)} \\
&\quad -x_{21} E_{(1,1,1,2,1,1)} - x_{22} E_{(1,1,1,2,2,1)} + x_{26} E_{(1,2,2,3,2,1)}
\end{aligned} \tag{4.17}$$

by (3.4),

$$\begin{aligned}
\iota(\eta_6) &= -\iota([E_{-\alpha_2}, \eta_4]) = -[\iota(E_{-\alpha_2}), \iota(\eta_4)] = -[E_{-\alpha_2}|_{\mathcal{A}} + E_{-\alpha_2}, \iota(\eta_4)] \\
&= P_6 - x_1 E'_{(0,1,0,1,1,1)} - x_2 E'_{(0,1,0,1,1)} - x_3 E'_{(0,1,0,1)} - x_4 E_{-\alpha_2} + x_7 E_{\alpha_3} \\
&\quad -x_6 \varrho(\sum_{i=2,4,5,6,7} \alpha_i) + x_9 E_{(0,0,1,1)} + x_{10} E_{(1,0,1)} + x_{12} E_{(1,0,1,1)} + x_{11} E_{(0,0,1,1,1)} \\
&\quad +x_{13} E_{(1,0,1,1,1)} + x_{14} E_{(0,0,1,1,1,1)} + x_{17} E_{(1,0,1,1,1,1)} - x_{20} E_{(1,1,2,2,1)} \\
&\quad -x_{23} E_{(1,1,2,2,1,1)} - x_{24} E_{(1,1,2,2,2,1)} - x_{25} E_{(1,1,2,3,2,1)}
\end{aligned} \tag{4.18}$$

by (3.3),

$$\begin{aligned}
\iota(\eta_7) &= -\iota([E_{-\alpha_2}, \eta_5]) = -[\iota(E_{-\alpha_2}), \iota(\eta_5)] = -[E_{-\alpha_2}|_{\mathcal{A}} + E_{-\alpha_2}, \iota(\eta_5)] \\
&= P_7 - x_1 E'_{(0,1,1,1,1,1)} - x_2 E'_{(0,1,1,1,1)} - x_3 E'_{(0,1,1,1)} - x_5 E_{-\alpha_2} - x_6 E_{-\alpha_3} \\
&\quad -x_7 \varrho(\sum_{i=2}^{7} \alpha_i) - x_9 E_{\alpha_4} + x_{10} E_{\alpha_1} + x_{15} E_{(1,0,1,1)} - x_{11} E_{(0,0,0,1,1)} \\
&\quad +x_{16} E_{(1,0,1,1,1)} - x_{14} E_{(0,0,0,1,1,1)} + x_{19} E_{(1,0,1,1,1,1)} + x_{20} E_{(1,1,1,2,1)} \\
&\quad +x_{23} E_{(1,1,1,2,1,1)} + x_{24} E_{(1,1,1,2,2,1)} + x_{26} E_{(1,1,2,3,2,1)}
\end{aligned} \tag{4.19}$$

by (3.3),

$$\begin{aligned}
\iota(\eta_8) &= -\iota([E_{-\alpha_1}, \eta_5]) = -[\iota(E_{-\alpha_1}), \iota(\eta_5)] = -[E_{-\alpha_1}|_{\mathcal{A}} + E_{-\alpha_1}, \iota(\eta_5)] \\
&= P_8 - x_1 E'_{(1,0,1,1,1,1)} - x_2 E'_{(1,0,1,1,1)} - x_3 E'_{(1,0,1,1)} - x_4 E'_{(1,0,1)} - x_5 E_{-\alpha_1} \\
&\quad -x_8 \varrho(\sum_{2\neq i\in\overline{1,7}} \alpha_i) + x_{10} E_{\alpha_2} + x_{12} E_{(0,1,0,1)} + x_{15} E_{(0,1,1,1)} + x_{13} E_{(0,1,0,1,1)} \\
&\quad +x_{16} E_{(0,1,1,1,1)} + x_{17} E_{(0,1,0,1,1,1)} + x_{19} E_{(0,1,1,1,1,1)} + x_{18} E_{(0,1,1,2,1)} \\
&\quad +x_{21} E_{(0,1,1,2,1,1)} + x_{22} E_{(0,1,1,2,2,1)} + x_{27} E_{(1,2,2,3,2,1)}
\end{aligned} \tag{4.20}$$

by (3.2),

$$\begin{aligned}
\iota(\eta_9) &= \iota([E_{-\alpha_4}, \eta_7]) = [\iota(E_{-\alpha_4}), \iota(\eta_7)] = [E_{-\alpha_4}|_{\mathcal{A}} + E_{-\alpha_4}, \iota(\eta_7)] \\
&= P_9 - x_1 E'_{(0,1,1,2,1,1)} - x_2 E'_{(0,1,1,2,1)} + x_4 E'_{(0,1,1,1)} - x_5 E'_{(0,1,0,1)} - x_6 E'_{(0,0,1,1)} \\
&\quad -x_9 \varrho(\alpha_4 + \sum_{i=2}^{7} \alpha_i) + x_7 E_{-\alpha_4} + x_{12} E_{\alpha_1} - x_{15} E_{(1,0,1)} - x_{11} E_{\alpha_5} \\
&\quad +x_{18} E_{(1,0,1,1,1)} - x_{14} E_{(0,0,0,0,1,1)} + x_{21} E_{(1,0,1,1,1,1)} - x_{20} E_{(1,1,1,1,1)} \\
&\quad -x_{23} E_{(1,1,1,1,1,1)} + x_{25} E_{(1,1,1,2,2,1)} - x_{26} E_{(1,1,2,2,2,1)}
\end{aligned} \tag{4.21}$$

by (3.5),

$$\begin{aligned}
\iota(\eta_{10}) &= -\iota([E_{-\alpha_1}, \eta_7]) = -[\iota(E_{-\alpha_1}), \iota(\eta_7)] = -[E_{-\alpha_1}|_{\mathcal{A}} + E_{-\alpha_1}, \iota(\eta_7)] \\
&= P_{10} - x_1 E'_{(1,1,1,1,1,1)} - x_2 E'_{(1,1,1,1,1)} - x_3 E'_{(1,1,1,1)} - x_8 E_{-\alpha_2} - x_6 E'_{(1,0,1)} \\
&\quad -x_{10}\varrho(\sum_{i=1}^{7} \alpha_i) - x_7 E_{-\alpha_1} - x_{12} E_{\alpha_4} - x_{15} E_{(0,0,1,1)} - x_{13} E_{(0,0,0,1,1)} \\
&\quad -x_{16} E_{(0,0,1,1,1)} - x_{17} E_{(0,0,0,1,1,1)} - x_{19} E_{(0,0,1,1,1,1)} - x_{20} E_{(0,1,1,2,1)} \\
&\quad -x_{23} E_{(0,1,1,2,1,1)} - x_{24} E_{(0,1,1,2,2,1)} + x_{27} E_{(1,1,2,3,2,1)} \qquad (4.22)
\end{aligned}$$

by (3.2),

$$\begin{aligned}
\iota(\eta_{11}) &= \iota([E_{-\alpha_5}, \eta_9]) = [\iota(E_{-\alpha_5}), \iota(\eta_9)] = [E_{-\alpha_5}|_{\mathcal{A}} + E_{-\alpha_5}, \iota(\eta_9)] \\
&= P_{11} - x_1 E'_{(0,1,1,2,2,1)} + x_3 E'_{(0,1,1,2,1)} + x_4 E'_{(0,1,1,1,1)} - x_5 E'_{(0,1,0,1,1)} \\
&\quad -x_{11}\varrho(\alpha_4 + \alpha_5 + \sum_{i=2}^{7} \alpha_i) - x_6 E'_{(0,0,1,1,1)} + x_7 E'_{(0,0,0,1,1)} + x_9 E_{-\alpha_5} \\
&\quad +x_{13} E_{\alpha_1} - x_{16} E_{(1,0,1)} - x_{18} E_{(1,0,1,1)} - x_{14} E_{\alpha_6} + x_{20} E_{(1,1,1,1)} \\
&\quad +x_{22} E_{(1,0,1,1,1,1)} - x_{24} E_{(1,1,1,1,1,1)} - x_{25} E_{(1,1,1,2,1,1)} + x_{26} E_{(1,1,2,2,1,1)} \qquad (4.23)
\end{aligned}$$

by (3.6),

$$\begin{aligned}
\iota(\eta_{12}) &= \iota([E_{-\alpha_4}, \eta_{10}]) = [\iota(E_{-\alpha_4}), \iota(\eta_{10})] = [E_{-\alpha_4}|_{\mathcal{A}} + E_{-\alpha_4}, \iota(\eta_{10})] \\
&= P_{12} - x_1 E'_{(1,1,1,2,1,1)} - x_2 E'_{(1,1,1,2,1)} + x_4 E'_{(1,1,1,1)} - x_6 E'_{(1,0,1,1)} \\
&\quad -x_{12}\varrho(\alpha_4 + \sum_{i=1}^{7} \alpha_i) - x_8 E'_{(0,1,0,1)} - x_9 E_{-\alpha_1} + x_{10} E_{-\alpha_4} \\
&\quad -x_{13} E_{\alpha_5} + x_{15} E_{\alpha_3} - x_{17} E_{(0,0,0,0,1,1)} - x_{18} E_{(0,0,1,1,1)} + x_{20} E_{(0,1,1,1,1)} \\
&\quad -x_{21} E_{(0,0,1,1,1,1)} + x_{23} E_{(0,1,1,1,1,1)} - x_{25} E_{(0,1,1,2,2,1)} - x_{27} E_{(1,1,2,2,2,1)} \qquad (4.24)
\end{aligned}$$

by (3.5),

$$\begin{aligned}
\iota(\eta_{13}) &= \iota([E_{-\alpha_5}, \eta_{12}]) = [\iota(E_{-\alpha_5}), \iota(\eta_{12})] = [E_{-\alpha_5}|_{\mathcal{A}} + E_{-\alpha_5}, \iota(\eta_{12})] \\
&= P_{13} - x_1 E'_{(1,1,1,2,2,1)} + x_3 E'_{(1,1,1,2,1)} + x_4 E'_{(1,1,1,1,1)} - x_8 E'_{(0,1,0,1,1)} \\
&\quad -x_{13}\varrho(\alpha_4 + \alpha_5 + \sum_{i=1}^{7} \alpha_i) - x_6 E'_{(1,0,1,1,1)} + x_{10} E'_{(0,0,0,1,1)} - x_{11} E_{-\alpha_1} \\
&\quad +x_{12} E_{-\alpha_5} + x_{16} E_{\alpha_3} - x_{17} E_{\alpha_6} + x_{18} E_{(0,0,1,1)} - x_{20} E_{(0,1,1,1)} \\
&\quad -x_{22} E_{(0,0,1,1,1,1)} + x_{24} E_{(0,1,1,1,1,1)} + x_{25} E_{(0,1,1,2,1,1)} + x_{27} E_{(1,1,2,2,1,1)} \qquad (4.25)
\end{aligned}$$

by (3.6),

$$\begin{aligned}
\iota(\eta_{14}) &= \iota([E_{-\alpha_6}, \eta_{11}]) = [\iota(E_{-\alpha_6}), \iota(\eta_{11})] = [E_{-\alpha_6}|_{\mathcal{A}} + E_{-\alpha_6}, \iota(\eta_{11})] \\
&= P_{14} + x_2 E'_{(0,1,1,2,2,1)} + x_3 E'_{(0,1,1,2,1,1)} + x_4 E'_{(0,1,1,1,1,1)} - x_5 E'_{(0,1,0,1,1,1)} \\
&\quad -x_{14}\varrho(\sum_{r=4}^{6} \alpha_r + \sum_{i=2}^{7} \alpha_i) - x_6 E'_{(0,0,1,1,1,1)} + x_7 E'_{(0,0,0,1,1,1)} + x_9 E'_{(0,0,0,0,1,1)} \\
&\quad +x_{11} E_{-\alpha_6} + x_{17} E_{\alpha_1} - x_{19} E_{(1,0,1)} - x_{21} E_{(1,0,1,1)} - x_{22} E_{(1,0,1,1,1)} \\
&\quad +x_{23} E_{(1,1,1,1)} + x_{24} E_{(1,1,1,1,1)} + x_{25} E_{(1,1,1,2,1)} - x_{26} E_{(1,1,2,2,1)} \qquad (4.26)
\end{aligned}$$

by (3.7),

$$\begin{aligned}
\iota(\eta_{15}) &= -\iota([E_{-\alpha_3},\eta_{12}]) = -[\iota(E_{-\alpha_2}),\iota(\eta_{12})] = -[E_{-\alpha_3}|_{\mathcal{A}} + E_{-\alpha_3},\iota(\eta_{12})] \\
&= P_{15} - x_1E'_{(1,1,2,2,1,1)} - x_2E'_{(1,1,2,2,1)} + x_5E'_{(1,1,1,1)} - x_7E'_{(1,0,1,1)} \\
&\quad -x_{15}\varrho(\alpha_3+\alpha_4+\sum_{i=1}^{7}\alpha_i) - x_8E'_{(0,1,1,1)} + x_9E'_{(1,0,1)} + x_{10}E'_{(0,0,1,1)} \\
&\quad -x_{12}E_{-\alpha_3} - x_{16}E_{\alpha_5} + x_{18}E_{(0,0,0,1,1)} - x_{19}E_{(0,0,0,0,1,1)} - x_{20}E_{(0,1,0,1,1)} \\
&\quad +x_{21}E_{(0,0,0,1,1,1)} - x_{23}E_{(0,1,0,1,1,1)} + x_{26}E_{(0,1,1,2,2,1)} + x_{27}E_{(1,1,1,2,2,1)} \qquad (4.27)
\end{aligned}$$

by (3.4),

$$\begin{aligned}
\iota(\eta_{16}) &= \iota([E_{-\alpha_5},\eta_{15}]) = [\iota(E_{-\alpha_5}),\iota(\eta_{15})] = [E_{-\alpha_5}|_{\mathcal{A}} + E_{-\alpha_5},\iota(\eta_{15})] \\
&= P_{16} - x_1E'_{(1,1,2,2,2,1)} + x_3E'_{(1,1,2,2,1)} + x_5E'_{(1,1,1,1,1)} - x_7E'_{(1,0,1,1,1)} \\
&\quad -x_{16}\varrho(\sum_{r=3}^{5}\alpha_r+\sum_{i=1}^{7}\alpha_i) - x_8E'_{(0,1,1,1,1)} + x_{10}E'_{(0,0,1,1,1)} + x_{11}E'_{(1,0,1)} \\
&\quad -x_{13}E_{-\alpha_3} + x_{15}E_{-\alpha_5} - x_{18}E_{\alpha_4} - x_{19}E_{\alpha_6} + x_{20}E_{(0,1,0,1)} + x_{22}E_{(0,0,0,1,1,1)} \\
&\quad -x_{24}E_{(0,1,0,1,1,1)} - x_{26}E_{(0,1,1,2,1,1)} - x_{27}E_{(1,1,1,2,1,1)} \qquad (4.28)
\end{aligned}$$

by (3.6),

$$\begin{aligned}
\iota(\eta_{17}) &= \iota([E_{-\alpha_6},\eta_{13}]) = [\iota(E_{-\alpha_6}),\iota(\eta_{13})] = [E_{-\alpha_6}|_{\mathcal{A}} + E_{-\alpha_6},\iota(\eta_{13})] \\
&= P_{17} + x_2E'_{(1,1,1,2,2,1)} + x_3E'_{(1,1,1,2,1,1)} + x_4E'_{(1,1,1,1,1,1)} - x_6E'_{(1,0,1,1,1,1)} \\
&\quad -x_{17}\varrho(\sum_{r=4}^{6}\alpha_r+\sum_{i=1}^{7}\alpha_i) - x_8E'_{(0,1,0,1,1,1)} + x_{10}E'_{(0,0,0,1,1,1)} - x_{14}E_{-\alpha_1} \\
&\quad +x_{12}E'_{(0,0,0,0,1,1)} + x_{13}E_{-\alpha_6} + x_{19}E_{\alpha_3} + x_{21}E_{(0,0,1,1)} - x_{23}E_{(0,1,1,1)} \\
&\quad +x_{22}E_{(0,0,1,1,1)} - x_{24}E_{(0,1,1,1,1)} - x_{25}E_{(0,1,1,2,1)} - x_{27}E_{(1,1,2,2,1)} \qquad (4.29)
\end{aligned}$$

by (3.7)

$$\begin{aligned}
\iota(\eta_{18}) &= \iota([E_{-\alpha_4},\eta_{16}]) = [\iota(E_{-\alpha_4}),\iota(\eta_{16})] = [E_{-\alpha_4}|_{\mathcal{A}} + E_{-\alpha_4},\iota(\eta_{16})] \\
&= P_{18} - x_1E'_{(1,1,2,3,2,1)} - x_4E'_{(1,1,2,2,1)} + x_5E'_{(1,1,1,2,1)} - x_8E'_{(0,1,1,2,1)} \\
&\quad -x_{18}\varrho(\alpha_4+\sum_{r=3}^{5}\alpha_r+\sum_{i=1}^{7}\alpha_i) - x_9E'_{(1,0,1,1,1)} + x_{11}E'_{(1,0,1,1)} + x_{12}E'_{(0,0,1,1,1)} \\
&\quad -x_{13}E'_{(0,0,1,1)} - x_{15}E'_{(0,0,0,1,1)} + x_{16}E_{-\alpha_4} - x_{21}E_{\alpha_6} - x_{20}E_{\alpha_2} + x_{22}E_{(0,0,0,0,1,1)} \\
&\quad -x_{25}E_{(0,1,0,1,1,1)} + x_{26}E_{(0,1,1,1,1,1)} + x_{27}E_{(1,1,1,1,1,1)} \qquad (4.30)
\end{aligned}$$

by (3.5),

$$\begin{aligned}
\iota(\eta_{19}) &= \iota([E_{-\alpha_6},\eta_{16}]) = [\iota(E_{-\alpha_6}),\iota(\eta_{16})] = [E_{-\alpha_6}|_{\mathcal{A}} + E_{-\alpha_6},\iota(\eta_{16})] \\
&= P_{19} + x_2E'_{(1,1,2,2,2,1)} + x_3E'_{(1,1,2,2,1,1)} + x_5E'_{(1,1,1,1,1,1)} - x_7E'_{(1,0,1,1,1,1)} \\
&\quad -x_{19}\varrho(\sum_{r=3}^{6}\alpha_r+\sum_{i=1}^{7}\alpha_i) - x_8E'_{(0,1,1,1,1,1)} + x_{10}E'_{(0,0,1,1,1,1)} + x_{14}E'_{(1,0,1)} \\
&\quad +x_{15}E'_{(0,0,0,0,1,1)} + x_{16}E_{-\alpha_6} - x_{17}E_{-\alpha_3} - x_{21}E_{\alpha_4} + x_{23}E_{(0,1,0,1)} \\
&\quad -x_{22}E_{(0,0,0,1,1)} + x_{24}E_{(0,1,0,1,1)} + x_{26}E_{(0,1,1,2,1)} + x_{27}E_{(1,1,1,2,1)} \qquad (4.31)
\end{aligned}$$

by (3.7),

$$\begin{aligned}
\iota(\eta_{20}) &= \iota([E_{-\alpha_2},\eta_{18}]) = [\iota(E_{-\alpha_2}),\iota(\eta_{18})] = [E_{-\alpha_2}|_{\mathcal{A}} + E_{-\alpha_2},\iota(\eta_{18})] \\
&= P_{20} - x_1E'_{(1,2,2,3,2,1)} + x_6E'_{(1,1,2,2,1)} - x_7E'_{(1,1,1,2,1)} + x_{10}E'_{(0,1,1,2,1)} \\
&\quad -x_{20}\varrho(\alpha_4 + \sum_{r=2}^{5}\alpha_r + \sum_{i=1}^{7}\alpha_i) + x_9E'_{(1,1,1,1,1)} - x_{11}E'_{(1,1,1,1)} - x_{12}E'_{(0,1,1,1,1)} \\
&\quad +x_{13}E'_{(0,1,1,1)} + x_{15}E'_{(0,1,0,1,1)} - x_{16}E_{(0,1,0,1)} + x_{18}E_{-\alpha_2} - x_{23}E_{\alpha_6} \\
&\quad +x_{24}E_{(0,0,0,0,1,1)} - x_{25}E_{(0,0,0,1,1,1)} + x_{26}E_{(0,0,1,1,1,1)} + x_{27}E_{(1,0,1,1,1,1)}
\end{aligned} \tag{4.32}$$

by (3.3),

$$\begin{aligned}
\iota(\eta_{21}) &= \iota([E_{-\alpha_4},\eta_{19}]) = [\iota(E_{-\alpha_4}),\iota(\eta_{19})] = [E_{-\alpha_4}|_{\mathcal{A}} + E_{-\alpha_4},\iota(\eta_{19})] \\
&= P_{21} + x_2E'_{(1,1,2,3,2,1)} - x_4E'_{(1,1,2,2,1,1)} + x_5E'_{(1,1,1,2,1,1)} - x_8E'_{(0,1,1,2,1,1)} \\
&\quad -x_{21}\varrho(\alpha_4 + \sum_{r=3}^{6}\alpha_r + \sum_{i=1}^{7}\alpha_i) - x_9E'_{(1,0,1,1,1,1)} + x_{12}E'_{(0,0,1,1,1,1)} + x_{14}E'_{(1,0,1,1)} \\
&\quad -x_{15}E'_{(0,0,0,1,1,1)} - x_{17}E'_{(0,0,1,1)} + x_{18}E_{-\alpha_6} + x_{19}E_{-\alpha_4} - x_{23}E_{\alpha_2} \\
&\quad -x_{22}E_{\alpha_5} + x_{25}E_{(0,1,0,1,1)} - x_{26}E_{(0,1,1,1,1)} - x_{27}E_{(1,1,1,1,1)}
\end{aligned} \tag{4.33}$$

by (3.5),

$$\begin{aligned}
\iota(\eta_{22}) &= \iota([E_{-\alpha_5},\eta_{21}]) = [\iota(E_{-\alpha_5}),\iota(\eta_{21})] = [E_{-\alpha_5}|_{\mathcal{A}} + E_{-\alpha_5},\iota(\eta_{21})] \\
&= P_{22} - x_3E'_{(1,1,2,3,2,1)} - x_4E'_{(1,1,2,2,2,1)} - x_{22}\varrho(\alpha_4 + \alpha_5 + \sum_{r=3}^{6}\alpha_r + \sum_{i=1}^{7}\alpha_i) \\
&\quad +x_5E'_{(1,1,1,2,2,1)} - x_8E'_{(0,1,1,2,2,1)} - x_{11}E'_{(1,0,1,1,1,1)} + x_{13}E'_{(0,0,1,1,1,1)} + x_{14}E'_{(1,0,1,1,1)} \\
&\quad -x_{16}E'_{(0,0,0,1,1,1)} - x_{17}E'_{(0,0,1,1.1)} - x_{18}E'_{(0,0,0,0,1,1)} + x_{19}E'_{(0,0,0,1,1)} \\
&\quad +x_{21}E_{-\alpha_5} - x_{24}E_{\alpha_2} - x_{25}E_{(0,1,0,1)} + x_{26}E_{(0,1,1,1)} + x_{27}E_{(1,1,1,1)}
\end{aligned} \tag{4.34}$$

by (3.6),

$$\begin{aligned}
\iota(\eta_{23}) &= \iota([E_{-\alpha_2},\eta_{21}]) = [\iota(E_{-\alpha_2}),\iota(\eta_{21})] = [E_{-\alpha_2}|_{\mathcal{A}} + E_{-\alpha_2},\iota(\eta_{21})] \\
&= P_{23} + x_2E'_{(1,2,2,3,2,1)} + x_6E'_{(1,1,2,2,1,1)} - x_7E'_{(1,1,1,2,1,1)} + x_9E'_{(1,1,1,1,1,1)} \\
&\quad -x_{23}\varrho(\alpha_4 + \sum_{r=2}^{6}\alpha_r + \sum_{i=1}^{7}\alpha_i) + x_{10}E'_{(0,1,1,2,1,1)} - x_{12}E'_{(0,1,1,1,1,1)} - x_{14}E'_{(1,1,1,1)} \\
&\quad +x_{15}E'_{(0,1,0,1,1,1)} + x_{17}E'_{(0,1,1,1)} + x_{20}E_{-\alpha_6} - x_{19}E_{(0,1,0,1)} + x_{21}E_{-\alpha_2} \\
&\quad -x_{24}E_{\alpha_5} + x_{25}E_{(0,0,0,1,1)} - x_{26}E_{(0,0,1,1,1)} - x_{27}E_{(1,0,1,1,1)}
\end{aligned} \tag{4.35}$$

by (3.3),

$$\begin{aligned}
\iota(\eta_{24}) &= \iota([E_{-\alpha_2},\eta_{22}]) = [\iota(E_{-\alpha_2}),\iota(\eta_{22})] = [E_{-\alpha_2}|_{\mathcal{A}} + E_{-\alpha_2},\iota(\eta_{22})] \\
&= P_{24} - x_3E'_{(1,2,2,3,2,1)} + x_6E'_{(1,1,2,2,2,1)} - x_{24}\varrho(\alpha_4 + \alpha_5 + \sum_{r=2}^{6}\alpha_r + \sum_{i=1}^{7}\alpha_i) \\
&\quad -x_7E'_{(1,1,1,2,2,1)} + x_{10}E'_{(0,1,1,2,2,1)} + x_{11}E'_{(1,1,1,1,1,1)} - x_{13}E'_{(0,1,1,1,1,1)} \\
&\quad -x_{14}E'_{(1,1,1,1,1)} + x_{16}E'_{(0,1,0,1,1,1)} + x_{17}E'_{(0,1,1,1,1)} - x_{19}E'_{(0,1,0,1,1)} - x_{20}E'_{(0,0,0,0,1,1)} \\
&\quad +x_{22}E_{-\alpha_2} + x_{23}E_{-\alpha_5} - x_{25}E_{\alpha_4} + x_{26}E_{(0,0,1,1)} + x_{27}E_{(1,0,1,1)}
\end{aligned} \tag{4.36}$$

by (3.3),

$$\begin{aligned}\iota(\eta_{25}) &= \iota([E_{-\alpha_4},\eta_{24}]) = [\iota(E_{-\alpha_4}),\iota(\eta_{24})] = [E_{-\alpha_4}|_{\mathcal{A}} + E_{-\alpha_4},\iota(\eta_{24})]\\ &= P_{25} + x_4E'_{(1,2,2,3,2,1)} + x_6E'_{(1,1,2,3,2,1)} - x_{25}\varrho(2\alpha_4+\alpha_5+\sum_{r=2}^{6}\alpha_r+\sum_{i=1}^{7}\alpha_i)\\ &\quad -x_9E'_{(1,1,1,2,2,1)} + x_{12}E'_{(0,1,1,2,2,1)} + x_{11}E'_{(1,1,1,2,1,1)} - x_{13}E'_{(0,1,1,2,1,1)}\\ &\quad -x_{14}E'_{(1,1,1,2,1)} + x_{17}E'_{(0,1,1,2,1)} + x_{18}E'_{(0,1,0,1,1,1)} + x_{20}E'_{(0,0,0,1,1,1)} - x_{21}E'_{(0,1,0,1,1)}\\ &\quad +x_{22}E'_{(0,1,0,1)} - x_{23}E'_{(0,0,0,1,1)} + x_{24}E_{-\alpha_4} - x_{26}E_{\alpha_3} - x_{27}E_{(1,0,1)} \qquad (4.37)\end{aligned}$$

by (3.5),

$$\begin{aligned}\iota(\eta_{26}) &= \iota([E_{-\alpha_3},\eta_{25}]) = [\iota(E_{-\alpha_3}),\iota(\eta_{25})] = [E_{-\alpha_3}|_{\mathcal{A}} + E_{-\alpha_3},\iota(\eta_{25})]\\ &= P_{26} - x_5E'_{(1,2,2,3,2,1)} - x_7E'_{(1,1,2,3,2,1)} - x_{26}\varrho(\alpha_4+\sum_{s=3}^{5}+\sum_{r=2}^{6}\alpha_r+\sum_{i=1}^{7}\alpha_i)\\ &\quad +x_9E'_{(1,1,2,2,2,1)} - x_{15}E'_{(0,1,1,2,2,1)} - x_{11}E'_{(1,1,2,2,1,1)} + x_{16}E'_{(0,1,1,2,1,1)}\\ &\quad +x_{14}E'_{(1,1,2,2,1)} - x_{19}E'_{(0,1,1,2,1)} - x_{18}E'_{(0,1,1,1,1,1)} - x_{20}E'_{(0,0,1,1,1,1)} + x_{21}E'_{(0,1,1,1,1)}\\ &\quad -x_{22}E'_{(0,1,1,1)} + x_{23}E'_{(0,0,1,1,1)} - x_{24}E'_{(0,0,1,1)} + x_{25}E_{-\alpha_3} + x_{27}E_{\alpha_1} \qquad (4.38)\end{aligned}$$

by (3.4),

$$\begin{aligned}\iota(\eta_{27}) &= -\iota([E_{-\alpha_1},\eta_{26}]) = -[\iota(E_{-\alpha_1}),\iota(\eta_{26})] = -[E_{-\alpha_1}|_{\mathcal{A}} + E_{-\alpha_1},\iota(\eta_{26})]\\ &= P_{27} - x_8E'_{(1,2,2,3,2,1)} - x_{10}E'_{(1,1,2,3,2,1)} - x_{27}\varrho(\alpha_4+\sum_{s=3}^{5}+\sum_{r=1}^{6}\alpha_r+\sum_{i=1}^{7}\alpha_i)\\ &\quad +x_{12}E'_{(1,1,2,2,2,1)} - x_{15}E'_{(1,1,1,2,2,1)} - x_{13}E'_{(1,1,2,2,1,1)} + x_{16}E'_{(1,1,1,2,1,1)}\\ &\quad +x_{17}E'_{(1,1,2,2,1)} - x_{19}E'_{(1,1,1,2,1)} - x_{18}E'_{(1,1,1,1,1,1)} - x_{20}E'_{(1,0,1,1,1,1)} + x_{21}E'_{(1,1,1,1,1)}\\ &\quad -x_{22}E'_{(1,1,1,1)} + x_{23}E'_{(1,0,1,1,1)} - x_{24}E'_{(1,0,1,1)} + x_{25}E'_{(1,0,1)} - x_{26}E_{-\alpha_1} \qquad (4.39)\end{aligned}$$

by (3.2). Note that the coefficient of x_i in the above $\iota(\eta_i)$ is exactly $\varrho(\beta)$ if $\eta_i = E_\beta$.

Recall $\mathcal{A} = \mathbb{C}[x_1,...,x_{27}]$. Let M be a $\mathcal{G}^{E_6}$-module and set

$$\widehat{M} = \mathcal{A}\otimes_{\mathbb{C}} M. \tag{4.40}$$

We identify

$$f\otimes v = fv \qquad \text{for } f\in\mathcal{A},\ v\in M. \tag{4.41}$$

Recall the Lie algebra $\widehat{\mathcal{K}}$ defined via (4.1)-(4.8). Fix $c\in\mathbb{C}$. Then $\widehat{M}$ becomes a $\widehat{\mathcal{K}}$-module with the action defined by

$$d(fw) = d(f)w,\ \ \kappa(fw) = cfw,\ \ (gu)(fw) = fgu(w) \tag{4.42}$$

for $d\in\mathcal{W}_{27},\ f,g\in\mathcal{A},\ w\in M$ and $u\in\mathcal{G}^{E_6}$.

Since the linear map $\iota:\mathcal{G}^{E_7}\to\widehat{\mathcal{K}}$ defined in (4.10)-(4.12) is a Lie algebra monomorphism, $\widehat{M}$ becomes a $\mathcal{G}^{E_7}$-module with the action defined by

$$\xi(w) = \iota(\xi)(w) \qquad \text{for } \xi\in\mathcal{G}^{E_7},\ w\in\widehat{M}. \tag{4.43}$$

In fact, we have:

Theorem 4.1. *The map $M \mapsto \widehat{M}$ gives a functor from $\mathcal{G}^{E_6}$-**Mod** to $\mathcal{G}^{E_7}$-**Mod**.*

We remark that the module $\widehat{M}$ is not a generalized module in general because it may not be equal to $U(\mathcal{G})(M) = U(\mathcal{G}_-)(M)$.

Proposition 4.2. *If M is an irreducible $\mathcal{G}^{E_6}$-module, then $U(\mathcal{G}_-)(M)$ is an irreducible $\mathcal{G}^{E_7}$-module.*

Proof. Note that for any $i \in \overline{1,27}$, $f \in \mathcal{A}$ and $v \in M$, (4.12), (4.42) and (4.43) imply

$$\xi_i(fv) = \partial_{x_i}(f)v. \tag{4.44}$$

Let W be any nonzero $\mathcal{G}^{E_7}$-submodule. The above expression shows that $W \bigcap M \neq \{0\}$. According to (4.42), $W \bigcap M$ is a $\mathcal{G}^{E_6}$-submodule. By the irreducibility of M, $M \subset W$. Thus $U(\mathcal{G}_-)(M) \subset W$. So $U(\mathcal{G}_-)(M) = W$ is irreducible. $\Box$

By the above proposition, the map $M \mapsto U(\mathcal{G}_-)(M)$ is a polynomial extension from irreducible $\mathcal{G}^{E_6}$-modules to irreducible $\mathcal{G}^{E_7}$-modules.

5 Irreducibility

In this section, we want to determine the irreducibility of $\mathcal{G}^{E_6}$-modules $\widehat{M}$.

Note that $\widehat{M}$ can be viewed as a $\mathcal{G}^{E_6}$-module. Indeed, (4.11) and (4.43) show

$$u(fv) = u(f)v + fu(v) \qquad \text{for } u \in \mathcal{G}^{E_6},\ f \in \mathcal{A},\ v \in M \tag{5.1}$$

(cf. (2.50)-(2.87)). So $\widehat{M} = \mathcal{A} \otimes_{\mathbb{C}} M$ is a tensor module of $\mathcal{G}^{E_6}$. Write

$$\eta^\alpha = \prod_{i=1}^{27} \eta_i^{\alpha_i}, \ \ |\alpha| = \sum_{i=1}^{27} \alpha_i \ \text{ for } \ \alpha = (\alpha_1, \alpha_2, ..., \alpha_{27}) \in \mathbb{N}^{27} \tag{5.2}$$

(cf. (2.34)-(2.39)). Recall the Lie subalgebras $\mathcal{G}_\pm$ and $\mathcal{G}_0$ of $\mathcal{G}^{E_6}$ defined in (2.41). For $k \in \mathbb{N}$, we set

$$\mathcal{A}_k = \text{Span}_{\mathbb{C}}\{x^\alpha \mid \alpha \in \mathbb{N}^{27}; |\alpha| = k\}, \ \ \widehat{M}_{\langle k \rangle} = \mathcal{A}_k M \tag{5.3}$$

(cf. (2.47), (4.41)) and

$$(U(\mathcal{G}_-)(M))_{\langle k \rangle} = \text{Span}_{\mathbb{C}}\{\eta^\alpha(M) \mid \alpha \in \mathbb{N}^{27},\ |\alpha| = k\}. \tag{5.4}$$

Moreover,

$$(U(\mathcal{G}_-)(M))_{\langle 0 \rangle} = \widehat{M}_{\langle 0 \rangle} = M. \tag{5.5}$$

Furthermore,

$$\widehat{M} = \bigoplus_{k=0}^{\infty} \widehat{M}_{\langle k \rangle}, \qquad U(\mathcal{G}_-)(M) = \bigoplus_{k=0}^{\infty} (U(\mathcal{G}_-)(M))_{\langle k \rangle}. \tag{5.6}$$

Next we define a linear transformation φ on $\widehat{M}$ determined by

$$\varphi(x^\alpha v) = \eta^\alpha(v) \qquad \text{for } \ \alpha \in \mathbb{N}^{27},\ v \in M. \tag{5.7}$$

Note that $\mathcal{A}_1 = \sum_{i=1}^{27} \mathbb{C}x_i$ forms the 27-dimensional $\mathcal{G}_0$-module of highest weight λ_6. According to (2.10) and (2.11), $\mathcal{G}_-$ forms a $\mathcal{G}_0$-module with respect to the adjoint representation, and the linear map from $\mathcal{A}_1$ to $\mathcal{G}_0$ determined by $x_i \mapsto \eta_i$ for $i \in \overline{1,27}$ gives a

$\mathcal{G}_0$-module isomorphism. Thus φ can also be viewed as a $\mathcal{G}_0$-module homomorphism from $\widehat{M}$ to $U(\mathcal{G}_0)(M)$. Moreover,

$$\varphi(\widehat{M}_{\langle k\rangle})=(U(\mathcal{G}_-)(M))_{\langle k\rangle}\qquad\text{for }\ k\in\mathbb{N}.\tag{5.8}$$

Observe that the fundamental weights of $\mathcal{G}^{E_6}$ are:

$$\lambda_1=\frac{1}{3}(4\alpha_1+3\alpha_2+5\alpha_3+6\alpha_4+4\alpha_5+2\alpha_6),\tag{5.9}$$

$$\lambda_2=\alpha_1+2\alpha_2+2\alpha_3+3\alpha_4+2\alpha_5+\alpha_6,\tag{5.10}$$

$$\lambda_3=\frac{1}{3}(5\alpha_1+6\alpha_2+10\alpha_3+12\alpha_4+8\alpha_5+4\alpha_6),\tag{5.11}$$

$$\lambda_4=2\alpha_1+3\alpha_2+4\alpha_3+6\alpha_4+4\alpha_5+2\alpha_6,\tag{5.12}$$

$$\lambda_5=\frac{1}{3}(4\alpha_1+6\alpha_2+8\alpha_3+12\alpha_4+10\alpha_5+5\alpha_6),\tag{5.13}$$

$$\lambda_6=\frac{1}{3}(2\alpha_1+3\alpha_2+4\alpha_3+6\alpha_4+5\alpha_5+4\alpha_6).\tag{5.14}$$

Using the above expressions, we can view $\lambda_i\in\mathcal{H}_{E_6}$ by (2.16). Then Casimir element of $\mathcal{G}^{E_6}$ is

$$\omega=\sum_{i=1}^{6}\lambda_i\alpha_i-\sum_{\beta\in\Phi_{E_6}}E_\beta E_{-\beta}\in U(\mathcal{G}^{E_6})\tag{5.15}$$

due to (2.12) and (2.13). The algebra $U(\mathcal{G}^{E_6})$ can be imbedded into the tensor algebra $U(\mathcal{G}^{E_6})\otimes U(\mathcal{G}^{E_6})$ by the associative algebra homomorphism $\delta: U(\mathcal{G}^{E_6})\to U(\mathcal{G}^{E_6})\otimes_{\mathbb{C}} U(\mathcal{G}^{E_6}))$ determined by

$$\delta(u)=u\otimes 1+1\otimes u\qquad\text{for }u\in\mathcal{G}^{E_6}.\tag{5.16}$$

Set

$$\tilde{\omega}=\frac{1}{2}(\delta(\omega)-\omega\otimes 1-1\otimes\omega)\in U(\mathcal{G}^{E_6})\otimes_{\mathbb{C}} U(\mathcal{G}^{E_6}).\tag{5.17}$$

Since $\sum_{i=1}^6\lambda_i\otimes\alpha_i$ is symmetric with respect to $\{\alpha_1,...,\alpha_6\}$ by (5.9)-(5.14),

$$\tilde{\omega}=\sum_{i=1}^{6}\lambda_i\otimes\alpha_i-\sum_{\beta\in\Phi_{E_6}}E_\beta\otimes E_{-\beta}.\tag{5.18}$$

Furthermore, $\tilde{\omega}$ acts on $\widehat{M}$ as a $\mathcal{G}^{E_6}$-module homomorphism via

$$(u_1\otimes u_2)(fw)=u_1(f)u_2(w)\qquad\text{for }\ u_1,u_2\in\mathcal{G}^{E_6},\ f\in\mathcal{A},\ w\in M.\tag{5.19}$$

Lemma 5.1. *We have* $\varphi|_{\widehat{M}_{\langle 1\rangle}}=(\tilde{\omega}-c)|_{\widehat{M}_{\langle 1\rangle}}$.

Proof. By (2.87) with Table 1 and (5.9)-(5.14)

$$\lambda_r|_{\mathcal{A}}=\sum_{i=1}^{27}c_{i,r}x_i\partial_{x_i}\qquad\text{for }\ r\in\overline{1,6}\tag{5.20}$$

with $c_{i,r}\in\mathbb{C}$, for instance,

$$c_{1,1}=\frac{2}{3},\ c_{1,2}=1,\ c_{1,3}=\frac{4}{3},\ c_{1,4}=2,\ c_{1,5}=\frac{5}{3},\ c_{1,6}=\frac{4}{3}.\tag{5.21}$$

On the other hand,

$$\alpha_7 = \frac{1}{3}[\widehat{\alpha} - (2\alpha_1 + 3\alpha_2 + 4\alpha_3 + 6\alpha_4 + 5\alpha_5 + 4\alpha_6)] \tag{5.22}$$

by (2.88). According to (4.10), (4.42), (5.19) and (5.21),

$$(\sum_{i=1}^{6} \lambda_i \otimes \alpha_i)(x_1 w) = c x_1 w - x_1 \alpha_7(w) \qquad \text{for } w \in M. \tag{5.23}$$

Thus (2.50)-(2.86) yield

$$\begin{aligned}\tilde{\omega}(x_1 w) \;=\; & [c x_1 + x_2 E_{\alpha_6} + x_3 E_{(0,0,0,0,1,1)} + x_4 E_{(0,0,0,1,1,1)} + x_5 E_{(0,0,1,1,1,1)} + x_6 E_{(0,1,0,1,1,1)} \\ & - x_1 \alpha_7(w) + x_7 E_{(0,1,1,1,1,1)} + x_8 E_{(1,0,1,1,1,1)} + x_9 E_{(0,1,1,2,1,1)} + x_{10} E_{(1,1,1,1,1,1)} \\ & + x_{11} E_{(0,1,1,2,2,1)} + x_{12} E_{(1,1,1,2,1,1)} + x_{13} E_{(1,1,1,2,2,1)} + x_{15} E_{(1,1,2,2,1,1)} \\ & + x_{16} E_{(1,1,2,2,2,1)} + x_{18} E_{(1,1,2,3,2,1)} + x_{20} E_{(1,2,2,3,2,1)}](w). \end{aligned} \tag{5.24}$$

Comparing (4.13) and (5.24), we get $\varphi(x_1 w) = (\tilde{\omega} - c)(x_1 w)$. According to (3.7), $-E_{-\alpha_6}(x_1) = x_2$. So for $v \in M$,

$$\begin{aligned} & \varphi(x_2 v) - \varphi(x_1 E_{-\alpha_6}(v)) = -(E_{-\alpha_6})(\varphi(x_1 v)) \\ = \;& -(E_{-\alpha_6})[(\tilde{\omega} - c)(x_1 v)] = (\tilde{\omega} - c)[-E_{-\alpha_6}(x_1 v)] \\ = \;& (\tilde{\omega} - c)[x_2 v - x_1 E_{-\alpha_6}(v)] \\ = \;& (\tilde{\omega} - c)(x_2 v) - (\tilde{\omega} - c)(x_1 E_{-\alpha_6})(v)) \\ = \;& (\tilde{\omega} - c)(x_2 v) - \varphi(x_1 E_{-\alpha_6}(v)), \end{aligned} \tag{5.25}$$

equivalently, $\varphi(x_2 v) = (\tilde{\omega} - c)(x_2 v)$. By (3.2)-(3.7), similar arguments as (5.25) and induction on i, we can prove

$$\varphi(x_i v) = (\tilde{\omega} - c)(x_i v) \qquad \text{for } \; i \in \overline{1,27}, \; v \in M, \tag{5.26}$$

that is, the lemma holds. $\qquad\square$

Recall that $\mathbb{N}$ is the set of nonnegative integers and the set of dominate integral weights is

$$\Lambda^+ = \sum_{r=1}^{6} \mathbb{N}\lambda_r. \tag{5.27}$$

Denote

$$\rho = \frac{1}{2} \sum_{\beta \in \Phi_{E_6}^+} \beta. \tag{5.28}$$

Then

$$\rho = \sum_{r=1}^{6} \lambda_r \tag{5.29}$$

(e.g., cf. [H]). For any $\mu \in \Lambda^+$, we denote by $V(\mu)$ the finite-dimensional irreducible $\mathcal{G}^{E_6}$-module with the highest weight μ and have

$$\omega|_{V(\mu)} = (\mu + 2\rho, \mu)\mathrm{Id}_{V(\mu)} \tag{5.30}$$

by (5.15). According to (2.87) and Table 1, the weight set of the $\mathcal{G}^{E_6}$-module $\mathcal{A}_1$ is

$$\Pi(\mathcal{A}_1) = \{\sum_{r=1}^{6} a_{i,r}\lambda_r \mid i \in \overline{1,27}\}. \tag{5.31}$$

Fixing $\lambda \in \Lambda^+$, we define

$$\Upsilon(\lambda) = \{\lambda + \mu \mid \mu \in \Pi(\mathcal{A}_1),\ \lambda + \mu \in \Lambda^+\}. \tag{5.32}$$

Lemma 5.2. *We have*:

$$\mathcal{A}_1 \otimes V(\lambda) \cong \bigoplus_{\lambda' \in \Upsilon(\lambda)} V(\lambda'). \tag{5.33}$$

Proof. Note that all the weight subspaces of $\mathcal{A}_1$ are one-dimensional. Thus all the irreducible components of $\mathcal{A}_1 \otimes V(\lambda)$ are of multiplicity one. Since

$$\rho + \lambda + \mu \in \Lambda^+ \qquad \text{for}\ \ \mu \in \Pi(\mathcal{A}_1), \tag{5.34}$$

the tensor theory of finite-dimensional irreducible modules over a finite-dimensional simple Lie algebra (e.g., cf. [H]) says that $V(\lambda')$ is a component of $\mathcal{A}_1 \otimes V(\lambda)$ if and only if $\lambda' \in \Upsilon(\lambda)$. $\quad\Box$

Recall

$$\text{the highest weight of } \mathcal{A}_1 = \lambda_6 \tag{5.35}$$

by Table 1. Thus the eigenvalues of $\tilde{\omega}|_{\widehat{V(\lambda)}_{\langle 1\rangle}}$ are

$$\{[(\lambda' + 2\rho, \lambda') - (\lambda + 2\rho, \lambda) - (\lambda_6 + 2\rho, \lambda_6)]/2 \mid \lambda' \in \Upsilon(\lambda)\} \tag{5.36}$$

by (5.17) and (5.19). Define

$$\ell_\omega(\lambda) = \min\{[(\lambda' + 2\rho, \lambda') - (\lambda + 2\rho, \lambda) - (\lambda_6 + 2\rho, \lambda_6)]/2 \mid \lambda' \in \Upsilon(\lambda)\}, \tag{5.37}$$

which will be used to determine the irreducibility of $\widehat{V(\lambda)}$. If $\lambda' = \lambda + \lambda_6 - \alpha \in \Upsilon(\lambda)$ with $\alpha \in \Phi^+_{D_6}$, then

$$(\lambda' + 2\rho, \lambda') - (\lambda + 2\rho, \lambda) - (\lambda_6 + 2\rho, \lambda_6) = 2[(\lambda, \lambda_6) + 1 - (\rho + \lambda + \lambda_6, \alpha)]. \tag{5.38}$$

Recall the differential operators $P_1, ..., P_{27}$ given in (3.58)-(3.102). We also view the elements of $\mathcal{A}$ as the multiplication operators on $\mathcal{A}$. Recall ζ_1 in (3.1). It turns out that we need the following lemma in order to determine the irreducibility of $\widehat{V(\lambda)}$.

Lemma 5.3. *As operators on* $\mathcal{A}$:

$$\begin{aligned} & P_{14}x_1 + P_1x_{14} + P_{11}x_2 + P_2x_{11} + P_9x_3 + P_3x_9 + P_7x_4 + P_4x_7 - P_6x_5 - P_5x_6 \\ = \ & \zeta_1(D-8) + \chi\partial_{x_{27}}. \end{aligned} \tag{5.39}$$

Proof. According to (3.58), (3.77)-(3.82), (3.84), (3.86) and (3.89), we find that

$$\begin{aligned} & P_{14}x_1 + P_1x_{14} + P_{11}x_2 + P_2x_{11} + P_9x_3 + P_3x_9 + P_7x_4 + P_4x_7 - P_6x_5 - P_5x_6 \\ = \ & -8\zeta_1 + x_1P_{14} + x_{14}P_1 + x_2P_{11} + x_{11}P_2 + x_3P_9 \\ & +x_9P_3 + x_4P_7 + x_7P_4 - x_5P_6 - x_6P_5 \end{aligned} \tag{5.40}$$

and

$$\begin{aligned}
& x_1P_{14}+x_{14}P_1+x_2P_{11}+x_{11}P_2+x_3P_9+x_9P_3+x_4P_7+x_7P_4-x_5P_6-x_6P_5 \\
={}& x_1(x_{14}D-\zeta_1\partial_{x_1}-\zeta_8\partial_{x_8}-\zeta_{10}\partial_{x_{10}}+\zeta_{12}\partial_{x_{12}}-\zeta_{13}\partial_{x_{13}}-\zeta_{15}\partial_{x_{15}}+\zeta_{16}\partial_{x_{16}}-\zeta_{18}\partial_{x_{18}} \\
& -\zeta_{20}\partial_{x_{20}}+\zeta_{27}\partial_{x_{27}})+x_{14}(x_1D-\zeta_1\partial_{x_{14}}-\zeta_2\partial_{x_{17}}-\zeta_3\partial_{x_{19}}+\zeta_4\partial_{x_{21}}-\zeta_5\partial_{x_{22}}-\zeta_6\partial_{x_{23}} \\
& +\zeta_7\partial_{x_{24}}-\zeta_9\partial_{x_{25}}-\zeta_{11}\partial_{x_{26}}-\zeta_{14}\partial_{x_{27}})+x_2(x_{11}D-\zeta_1\partial_{x_2}+\zeta_5\partial_{x_8}+\zeta_7\partial_{x_{10}}-\zeta_9\partial_{x_{12}} \\
& +\zeta_{11}\partial_{x_{15}}+\zeta_{13}\partial_{x_{17}}-\zeta_{16}\partial_{x_{19}}+\zeta_{18}\partial_{x_{21}}+\zeta_{20}\partial_{x_{23}}+\zeta_{26}\partial_{x_{27}})+x_{11}(x_2D-\zeta_1\partial_{x_{11}} \\
& -\zeta_2\partial_{x_{13}}-\zeta_3\partial_{x_{16}}+\zeta_4\partial_{x_{18}}-\zeta_6\partial_{x_{20}}-\zeta_8\partial_{x_{22}}+\zeta_{10}\partial_{x_{24}}-\zeta_{12}\partial_{x_{25}}-\zeta_{15}\partial_{x_{26}}+\zeta_{17}\partial_{x_{27}}) \\
& +x_3(x_9D-\zeta_1\partial_{x_3}-\zeta_4\partial_{x_8}-\zeta_6\partial_{x_{10}}+\zeta_9\partial_{x_{13}}-\zeta_{12}\partial_{x_{17}}-\zeta_{11}\partial_{x_{16}}+\zeta_{15}\partial_{x_{19}}-\zeta_{18}\partial_{x_{22}} \\
& -\zeta_{20}\partial_{x_{24}}+\zeta_{25}\partial_{x_{27}})+x_9(x_3D-\zeta_1\partial_{x_9}-\zeta_2\partial_{x_{12}}-\zeta_3\partial_{x_{15}}+\zeta_5\partial_{x_{18}}-\zeta_7\partial_{x_{20}}-\zeta_8\partial_{x_{21}} \\
& +\zeta_{10}\partial_{x_{23}}-\zeta_{13}\partial_{x_{25}}-\zeta_{16}\partial_{x_{26}}+\zeta_{19}\partial_{x_{27}})+x_4(x_7D-\zeta_1\partial_{x_4}+\zeta_3\partial_{x_8}+\zeta_6\partial_{x_{12}}-\zeta_7\partial_{x_{13}} \\
& +\zeta_{10}\partial_{x_{17}}+\zeta_{11}\partial_{x_{18}}-\zeta_{15}\partial_{x_{21}}+\zeta_{16}\partial_{x_{22}}+\zeta_{20}\partial_{x_{25}}+\zeta_{24}\partial_{x_{27}})+x_7(x_4D-\zeta_1\partial_{x_7} \\
& -\zeta_2\partial_{x_{10}}-\zeta_4\partial_{x_{15}}+\zeta_5\partial_{x_{16}}-\zeta_8\partial_{x_{19}}-\zeta_9\partial_{x_{20}}+\zeta_{12}\partial_{x_{23}}-\zeta_{13}\partial_{x_{24}}-\zeta_{18}\partial_{x_{26}}+\zeta_{21}\partial_{x_{27}}) \\
& -x_5(x_6D+\zeta_1\partial_{x_5}+\zeta_2\partial_{x_8}-\zeta_6\partial_{x_{15}}+\zeta_7\partial_{x_{16}}-\zeta_9\partial_{x_{18}}-\zeta_{10}\partial_{x_{19}}+\zeta_{12}\partial_{x_{21}} \\
& -\zeta_{13}\partial_{x_{22}}+\zeta_{20}\partial_{x_{26}}-\zeta_{23}\partial_{x_{27}})-x_6(x_5D+\zeta_1\partial_{x_6}-\zeta_3\partial_{x_{10}}+\zeta_4\partial_{x_{12}} \\
& -\zeta_5\partial_{x_{13}}+\zeta_8\partial_{x_{17}}+\zeta_{11}\partial_{x_{20}}-\zeta_{15}\partial_{x_{23}}+\zeta_{16}\partial_{x_{24}}-\zeta_{18}\partial_{x_{25}}-\zeta_{22}\partial_{x_{27}}) \\
={}& \zeta_1(2D-\sum_{i=1}^{7}x_i\partial_{x_i}-x_9\partial_{x_9}-x_{11}\partial_{x_{11}}-x_{14}\partial_{x_{14}})-(x_1\zeta_8-x_2\zeta_5+x_3\zeta_4-x_4\zeta_3 \\
& +x_5\zeta_2)\partial_{x_8}-(x_1\zeta_{10}-x_2\zeta_7+x_3\zeta_6+x_7\zeta_2-x_6\zeta_3)\partial_{x_{10}}+(x_1\zeta_{12}-x_2\zeta_9-x_9\zeta_2 \\
& +x_4\zeta_6-x_6\zeta_4)\partial_{x_{12}}-(x_1\zeta_{13}+x_{11}\zeta_2-x_3\zeta_9+x_4\zeta_7-x_6\zeta_5)\partial_{x_{13}}-(x_1\zeta_{15}-x_2\zeta_{11} \\
& +x_9\zeta_3+x_7\zeta_4-x_5\zeta_6)\partial_{x_{15}}+(x_1\zeta_{16}-x_{11}\zeta_3-x_3\zeta_{11}+x_7\zeta_5-x_5\zeta_7)\partial_{x_{16}}-(x_{14}\zeta_2 \\
& -x_2\zeta_{13}+x_3\zeta_{12}-x_4\zeta_{10}+x_6\zeta_8)\partial_{x_{17}}-(x_1\zeta_{18}-x_{11}\zeta_4-x_9\zeta_5-x_4\zeta_{11}-x_5\zeta_9)\partial_{x_{18}} \\
& -(x_{14}\zeta_3+x_2\zeta_{16}-x_3\zeta_{15}+x_7\zeta_8-x_5\zeta_{10})\partial_{x_{19}}-(x_1\zeta_{20}+x_{11}\zeta_6+x_9\zeta_7+x_7\zeta_9 \\
& +x_6\zeta_{11})\partial_{x_{20}}+(x_{14}\zeta_4+x_2\zeta_{18}-x_9\zeta_8-x_4\zeta_{15}-x_5\zeta_{12})\partial_{x_{21}}-(x_{14}\zeta_5+x_{11}\zeta_8 \\
& +x_3\zeta_{18}-x_4\zeta_{16}-x_5\zeta_{13})\partial_{x_{22}}-(x_{14}\zeta_6-x_2\zeta_{20}-x_9\zeta_{10}-x_7\zeta_{12}-x_6\zeta_{15})\partial_{x_{23}} \\
& +(x_{14}\zeta_7+x_{11}\zeta_{10}-x_3\zeta_{20}-x_7\zeta_{13}-x_6\zeta_{16})\partial_{x_{24}}-(x_{14}\zeta_9+x_{11}\zeta_{12}+x_9\zeta_{13} \\
& -x_4\zeta_{20}-x_6\zeta_{18})\partial_{x_{25}}-(x_{14}\zeta_{11}+x_{11}\zeta_{15}+x_9\zeta_{16}+x_7\zeta_{18}+x_5\zeta_{20})\partial_{x_{26}}+(x_1\zeta_{27} \\
& -x_{14}\zeta_{14}+x_2\zeta_{26}+x_{11}\zeta_{17}+x_3\zeta_{25}+x_9\zeta_{19}+x_4\zeta_{24}+x_7\zeta_{21}+x_5\zeta_{23}+x_6\zeta_{22})\partial_{x_{27}} \\
={}& \zeta_1D+\chi\partial_{x_{27}}
\end{aligned}\tag{5.41}$$

because

$$x_1\zeta_8-x_2\zeta_5+x_3\zeta_4-x_4\zeta_3+x_5\zeta_2=\zeta_1x_8\tag{5.42}$$

by (3.1), (3.8)-(3.11) and (3.14),

$$x_1\zeta_{10}-x_2\zeta_7+x_3\zeta_6+x_7\zeta_2-x_6\zeta_3=\zeta_1x_{10}\tag{5.43}$$

by (3.1), (3.8), (3.9), (3.12), (3.13) and (3.16),

$$x_1\zeta_{12}-x_2\zeta_9-x_9\zeta_2+x_4\zeta_6-x_6\zeta_4=-\zeta_1x_{12}\tag{5.44}$$

by (3.1), (3.8), (3.10), (3.12), (3.15) and (3.18),

$$x_1\zeta_{13}+x_{11}\zeta_2-x_3\zeta_9+x_4\zeta_7-x_6\zeta_5=\zeta_1x_{13}\tag{5.45}$$

by (3.1), (3.8), (3.11), (3.13), (3.15) and (3.19),

$$x_1\zeta_{15} - x_2\zeta_{11} + x_9\zeta_3 + x_7\zeta_4 - x_5\zeta_6 = \zeta_1 x_{15} \tag{5.46}$$

by (3.1), (3.9), (3.10), (3.12), (3.17) and (3.21),

$$x_1\zeta_{16} - x_{11}\zeta_3 - x_3\zeta_{11} + x_7\zeta_5 - x_5\zeta_7 = -\zeta_1 x_{16} \tag{5.47}$$

by (3.1), (3.9), (3.11), (3.13), (3.17) and (3.22),

$$x_{14}\zeta_2 - x_2\zeta_{13} + x_3\zeta_{12} - x_4\zeta_{10} + x_6\zeta_8 = \zeta_1 x_{17} \tag{5.48}$$

by (3.1), (3.8), (3.14), (3.16), (3.18) and (3.19),

$$x_1\zeta_{18} - x_{11}\zeta_4 - x_9\zeta_5 - x_4\zeta_{11} - x_5\zeta_9 = \zeta_1 x_{18} \tag{5.49}$$

by (3.1), (3.10), (3.11), (3.15), (3.17) and (3.24),

$$x_{14}\zeta_3 + x_2\zeta_{16} - x_3\zeta_{15} + x_7\zeta_8 - x_5\zeta_{10} = \zeta_1 x_{19} \tag{5.50}$$

by (3.1), (3.9), (3.14), (3.16), (3.21) and (3.22),

$$x_1\zeta_{20} + x_{11}\zeta_6 + x_9\zeta_7 + x_7\zeta_9 + x_6\zeta_{11} = \zeta_1 x_{20} \tag{5.51}$$

by (3.1), (3.12), (3.13), (3.15), (3.17) and (3.26),

$$x_{14}\zeta_4 + x_2\zeta_{18} - x_9\zeta_8 - x_4\zeta_{15} - x_5\zeta_{12} = -\zeta_1 x_{21} \tag{5.52}$$

by (3.1), (3.10), (3.14), (3.18), (3.21) and (3.24),

$$x_{14}\zeta_5 + x_{11}\zeta_8 + x_3\zeta_{18} - x_4\zeta_{16} - x_5\zeta_{13} = \zeta_1 x_{22} \tag{5.53}$$

by (3.1), (3.11), (3.14), (3.19), (3.22) and (3.24),

$$x_{14}\zeta_6 - x_2\zeta_{20} - x_9\zeta_{10} - x_7\zeta_{12} - x_6\zeta_{15} = \zeta_1 x_{23} \tag{5.54}$$

by (3.1), (3.12), (3.16), (3.18), (3.21) and (3.26),

$$x_{14}\zeta_7 + x_{11}\zeta_{10} - x_3\zeta_{20} - x_7\zeta_{13} - x_6\zeta_{16} = -\zeta_1 x_{24} \tag{5.55}$$

by (3.1), (3.13), (3.16), (3.19), (3.22) and (3.26),

$$x_{14}\zeta_9 + x_{11}\zeta_{12} + x_9\zeta_{13} - x_4\zeta_{20} - x_6\zeta_{18} = \zeta_1 x_{25} \tag{5.56}$$

by (3.1), (3.15), (3.18), (3.19), (3.24) and (3.26),

$$x_{14}\zeta_{11} + x_{11}\zeta_{15} + x_9\zeta_{16} + x_7\zeta_{18} + x_5\zeta_{20} = \zeta_1 x_{26} \tag{5.57}$$

by (3.1), (3.17), (3.21), (3.22), (3.24) and (3.26),

$$\begin{aligned} & x_1\zeta_{27} - x_{14}\zeta_{14} + x_2\zeta_{26} + x_{11}\zeta_{17} + x_3\zeta_{25} + x_9\zeta_{19} + x_4\zeta_{24} \\ & + x_7\zeta_{21} + x_5\zeta_{23} + x_6\zeta_{22} = -\zeta_1 x_{27} + \chi \end{aligned} \tag{5.58}$$

by (3.1), (3.20), (3.23), (3.25), (3.27)-(3.33) and (3.137). □

Lemma 5.4. *As operators on* $\mathcal{A}$,

$$\sum_{14\neq i\in\overline{1,26}} P_i\zeta_{28-i} - P_{14}\zeta_{14} - P_{27}\zeta_1 = \chi(24-5D). \tag{5.59}$$

Proof. We calculate it by (3.1), (3.8)-(3.33), (3.58) and (3.77)-(3.102). In particular,

$$2(\zeta_1\zeta_{14} - \zeta_2\zeta_{11} - \zeta_3\zeta_9 + \zeta_4\zeta_7 - \zeta_5\zeta_6) = -2\chi x_1, \tag{5.60}$$

which is the coefficient of ∂_{x_1} in addition to the term containing D. According to (2.138), the operator on the left hand side of (5.59) is a $\mathcal{G}^{E_6}$-invariant differential operator. By symmetry,

$$\text{the coefficient of } \partial_{x_i} = -2\chi x_i \ \text{ for } \ i\in\overline{1,27}. \qquad \Box \tag{5.61}$$

We define the multiplication

$$f(gv) = (fg)v \qquad \text{for } \ f,g\in\mathcal{A},\ v\in M. \tag{5.62}$$

Furthermore, we have:

Lemma 5.5. *As operators on* $\widehat{M}$,

$$[(\sum_{14\neq i\in\overline{1,26}} \iota(\eta_i)\zeta_{28-i} - \iota(\eta_{14})\zeta_{14} - \iota(\eta_{27})\zeta_1]|_{\widehat{M}} = \chi(24-5D+3c). \tag{5.63}$$

Proof. By (4.13)-(4.39), the coefficient of E_{α_6} is

$$x_2\zeta_{27} - x_{14}\zeta_{17} - x_{17}\zeta_{15} - x_{19}\zeta_{12} - x_{21}\zeta_{10} - x_{23}\zeta_8. \tag{5.64}$$

Moreover, we use (3.14), (3.16), (3.18), (3.21), (3.23) and (3.33) to find that (5.64) is equal to zero. Since the left hand side of (5.63) is a $\mathcal{G}^{E_6}$-invariant differential operator, it is invariant under the action of the E_6 Weyl group. The transitivity the Weyl group on Φ_{E_6} yields that

$$\text{the coefficient of } E_\beta = 0 \ \text{ for any } \beta\in\Phi_{E_6}. \tag{5.65}$$

By (4.10), (4.42), (5.22) and Lemma 5.4,

$$\sum_{14\neq i\in\overline{1,26}} \iota(\eta_i)\zeta_{28-i} - \iota(\eta_{14})\zeta_{14} - \iota(\eta_{27})\zeta_1 = \chi(24-5D+3c) + \sum_{r=1}^{6} f_r\alpha_r \tag{5.66}$$

as the elements of $\widehat{\mathcal{K}}$ acting on $\widehat{M}$ (cf. (4.7)). The $\mathcal{G}^{E_6}$-invariancy implies

$$[E_\beta|_{\widehat{M}}, \sum_{r=1}^{6} f_r\alpha_r] = 0 \ \text{ for any } \beta\in\Phi_{E_6}. \tag{5.68}$$

Thus $\sum_{r=1}^6 f_r\alpha_r = 0$, that is, (3.63) holds. $\qquad\Box$

Next we calculate

$$\begin{aligned} T_1 \ = \ & \iota(\eta_{14})x_1 + \iota(\eta_1)x_{14} + \iota(\eta_{11})x_2 + \iota(\eta_2)x_{11} + \iota(\eta_9)x_3 \\ & +\iota(\eta_3)x_9 + \iota(\eta_7)x_4 + \iota(\eta_4)x_7 - \iota(\eta_6)x_5 - \iota(\eta_5)x_6 \end{aligned}$$

$$\begin{aligned} &= \zeta_1(D-8)+\chi\partial_{x_{27}}+\zeta_2E_{\alpha_1}-\zeta_3E_{(1,0,1)}+\zeta_4E_{(1,0,1,1)}-\zeta_5E_{(1,0,1,1,1)}+\zeta_6E_{(1,1,1,1)} \\ &\quad -\zeta_7E_{(1,1,1,1,1)}+\zeta_8E_{(1,0,1,1,1,1)}+\zeta_9E_{(1,1,1,2,1)}+\zeta_{10}E_{(1,1,1,1,1,1)}-\zeta_{11}E_{(1,1,2,2,1)} \\ &\quad -\zeta_{12}E_{(1,1,1,2,1,1)}+\zeta_{13}E_{(1,1,1,2,2,1)}+\zeta_{15}E_{(1,1,2,2,1,1)}-\zeta_{16}E_{(1,1,2,2,2,1)}+\zeta_{18}E_{(1,1,2,3,2,1)} \\ &\quad +\zeta_{20}E_{(1,2,2,3,2,1)}-2\zeta_1c+\frac{\zeta_1}{3}(4\alpha_1+3\alpha_2+5\alpha_3+6\alpha_4+4\alpha_5+2\alpha_6) \end{aligned} \tag{5.69}$$

by (3.1), (3.8)-(3.19), (3.21), (3.22), (3.24), (3.26), (4.13)-(4.19), (4.21), (4.23), (4.26), (5.20) and Lemma 5.3.

We define a $\mathcal{G}^{E_6}$-module structure on the space $\mathrm{End}\,\widehat{M}$ of linear transformations on $\widehat{M}$ by

$$\iota(u)(T)=[\iota(u),T]=\iota(u)T-T\iota(u)\qquad\text{for }\ u\in\mathcal{G}^{E_6},\ T\in\mathrm{End}\,\widehat{M} \tag{5.70}$$

(cf. (5.1)). It can be verified that T_1 is a $\mathcal{G}^{E_6}$-singular vector with weight λ_1 in $\mathrm{End}\,\widehat{M}$. So it generates the 27-dimensional module of highest weight λ_1. We set

$$\begin{aligned} T_2 &= -[\iota(E_{-\alpha_1}),T_1]=\zeta_2(D-8)-\chi\partial_{x_{26}}-\zeta_1E_{-\alpha_1}+\zeta_3E_{\alpha_3}-\zeta_4E_{(0,0,1,1)} \\ &\quad +\zeta_5E_{(0,0,1,1,1)}-\zeta_6E_{(0,1,1,1)}+\zeta_7E_{(0,1,1,1,1)}-\zeta_8E_{(0,0,1,1,1,1)}-\zeta_9E_{(0,1,1,2,1)} \\ &\quad -\zeta_{10}E_{(0,1,1,1,1,1)}-\zeta_{14}E_{(1,1,2,2,1)}+\zeta_{12}E_{(0,1,1,2,1,1)}-\zeta_{13}E_{(0,1,1,2,2,1)} \\ &\quad -\zeta_{17}E_{(1,1,2,2,1,1)}+\zeta_{19}E_{(1,1,2,2,2,1)}-\zeta_{21}E_{(1,1,2,3,2,1)}-\zeta_{23}E_{(1,2,2,3,2,1)} \\ &\quad -2\zeta_2c+\frac{\zeta_2}{3}(\alpha_1+3\alpha_2+5\alpha_3+6\alpha_4+4\alpha_5+2\alpha_6) \end{aligned} \tag{5.71}$$

by (2.10), (2.11), (3.2) and (3.41),

$$\begin{aligned} T_3 &= -[\iota(E_{-\alpha_3}),T_2]=\zeta_3(D-8)-\chi\partial_{x_{25}}+\zeta_1E'_{(1,0,1)}-\zeta_2E_{-\alpha_3}+\zeta_4E_{\alpha_4} \\ &\quad -\zeta_5E_{(0,0,0,1,1)}+\zeta_6E_{(0,1,0,1)}-\zeta_7E_{(0,1,0,1,1)}+\zeta_8E_{(0,0,0,1,1,1)}+\zeta_{11}E_{(0,1,1,2,1)} \\ &\quad +\zeta_{10}E_{(0,1,0,1,1,1)}+\zeta_{14}E_{(1,1,1,2,1)}-\zeta_{15}E_{(0,1,1,2,1,1)}+\zeta_{16}E_{(0,1,1,2,2,1)} \\ &\quad +\zeta_{17}E_{(1,1,1,2,1,1)}-\zeta_{19}E_{(1,1,1,2,2,1)}+\zeta_{22}E_{(1,1,2,3,2,1)}+\zeta_{24}E_{(1,2,2,3,2,1)} \\ &\quad -2\zeta_3c+\frac{\zeta_3}{3}(\alpha_1+3\alpha_2+2\alpha_3+6\alpha_4+4\alpha_5+2\alpha_6) \end{aligned} \tag{5.72}$$

by (2.10), (2.11), (3.4) and (3.43),

$$\begin{aligned} T_4 &= -[\iota(E_{-\alpha_4}),T_3]=\zeta_4(D-8)-\chi\partial_{x_{24}}-\zeta_1E'_{(1,0,1,1)}+\zeta_2E'_{(0,0,1,1)}-\zeta_3E_{-\alpha_4} \\ &\quad +\zeta_5E_{\alpha_5}+\zeta_6E_{\alpha_2}-\zeta_9E_{(0,1,0,1,1)}-\zeta_8E_{(0,0,0,0,1,1)}+\zeta_{11}E_{(0,1,1,1,1)} \\ &\quad +\zeta_{12}E_{(0,1,0,1,1,1)}+\zeta_{14}E_{(1,1,1,1,1)}-\zeta_{15}E_{(0,1,1,1,1,1)}+\zeta_{18}E_{(0,1,1,2,2,1)} \\ &\quad +\zeta_{17}E_{(1,1,1,1,1,1)}-\zeta_{21}E_{(1,1,1,2,2,1)}+\zeta_{22}E_{(1,1,2,2,2,1)}-\zeta_{25}E_{(1,2,2,3,2,1)} \\ &\quad -2\zeta_4c+\frac{\zeta_4}{3}(\alpha_1+3\alpha_2+2\alpha_3+3\alpha_4+4\alpha_5+2\alpha_6) \end{aligned} \tag{5.73}$$

by (2.10), (2.11), (3.5) and (3.44),

$$\begin{aligned} T_5 &= -[\iota(E_{-\alpha_5}),T_4]=\zeta_5(D-8)-\chi\partial_{x_{23}}+\zeta_1E'_{(1,0,1,1,1)}-\zeta_2E'_{(0,0,1,1,1)} \\ &\quad +\zeta_3E'_{(0,0,0,1,1)}-\zeta_4E_{-\alpha_5}+\zeta_7E_{\alpha_2}-\zeta_9E_{(0,1,0,1)}+\zeta_8E_{\alpha_6}+\zeta_{11}E_{(0,1,1,1)} \\ &\quad +\zeta_{13}E_{(0,1,0,1,1,1)}+\zeta_{14}E_{(1,1,1,1)}-\zeta_{16}E_{(0,1,1,1,1,1)}+\zeta_{18}E_{(0,1,1,2,1,1)} \\ &\quad +\zeta_{19}E_{(1,1,1,1,1,1)}-\zeta_{21}E_{(1,1,1,2,1,1)}+\zeta_{22}E_{(1,1,2,2,1,1)}+\zeta_{26}E_{(1,2,2,3,2,1)} \\ &\quad -2\zeta_5c+\frac{\zeta_5}{3}(\alpha_1+3\alpha_2+2\alpha_3+3\alpha_4+\alpha_5+2\alpha_6) \end{aligned} \tag{5.74}$$

by (2.10), (2.11), (3.6) and (3.45),

$$\begin{aligned} T_6 &= -[\iota(E_{-\alpha_2}), T_4] = \zeta_6(D-8) - \chi\partial_{x_{22}} - \zeta_1 E'_{(1,1,1,1)} + \zeta_2 E'_{(0,1,1,1)} - \zeta_3 E'_{(0,1,0,1)} \\ &\quad + \zeta_7 E_{\alpha_5} - \zeta_4 E_{-\alpha_2} + \zeta_9 E_{(0,0,0,1,1)} - \zeta_{10} E_{(0,0,0,0,1,1)} - \zeta_{11} E_{(0,0,1,1,1)} \\ &\quad - \zeta_{12} E_{(0,0,0,1,1,1)} - \zeta_{14} E_{(1,0,1,1,1)} + \zeta_{15} E_{(0,0,1,1,1,1)} - \zeta_{20} E_{(0,1,1,2,2,1)} \\ &\quad - \zeta_{17} E_{(1,0,1,1,1,1)} + \zeta_{23} E_{(1,1,1,2,2,1)} - \zeta_{24} E_{(1,1,2,2,2,1)} - \zeta_{25} E_{(1,1,2,3,2,1)} \\ &\quad - 2\zeta_6 c + \frac{\zeta_6}{3}(\alpha_1 + 2\alpha_3 + 3\alpha_4 + 4\alpha_5 + 2\alpha_6) \end{aligned} \tag{5.75}$$

by (2.10), (2.11), (3.3) and (3.42),

$$\begin{aligned} T_7 &= -[\iota(E_{-\alpha_5}), T_6] = \zeta_7(D-8) - \chi\partial_{x_{21}} + \zeta_1 E'_{(1,1,1,1,1)} - \zeta_2 E'_{(0,1,1,1,1)} \\ &\quad + \zeta_3 E'_{(0,1,0,1,1)} - \zeta_6 E_{-\alpha_5} - \zeta_5 E_{-\alpha_2} + \zeta_9 E_{\alpha_4} + \zeta_{10} E_{\alpha_6} - \zeta_{11} E_{(0,0,1,1)} \\ &\quad - \zeta_{13} E_{(0,0,0,1,1,1)} - \zeta_{14} E_{(1,0,1,1)} + \zeta_{16} E_{(0,0,1,1,1,1)} - \zeta_{20} E_{(0,1,1,2,1,1)} \\ &\quad - \zeta_{19} E_{(1,0,1,1,1,1)} + \zeta_{23} E_{(1,1,1,2,1,1)} - \zeta_{24} E_{(1,1,2,2,1,1)} + \zeta_{26} E_{(1,1,2,3,2,1)} \\ &\quad - 2\zeta_7 c + \frac{\zeta_7}{3}(\alpha_1 + 2\alpha_3 + 3\alpha_4 + \alpha_5 + 2\alpha_6) \end{aligned} \tag{5.76}$$

by (2.10), (2.11), (3.6) and (3.45),

$$\begin{aligned} T_8 &= -[\iota(E_{-\alpha_6}), T_5] = \zeta_8(D-8) - \chi\partial_{x_{20}} - \zeta_1 E'_{(1,0,1,1,1,1)} + \zeta_2 E'_{(0,0,1,1,1,1)} \\ &\quad - \zeta_3 E'_{(0,0,0,1,1,1)} + \zeta_4 E'_{(0,0,0,0,1,1)} - \zeta_5 E_{-\alpha_6} + \zeta_{10} E_{\alpha_2} - \zeta_{12} E_{(0,1,0,1)} \\ &\quad + \zeta_{13} E_{(0,1,0,1,1)} + \zeta_{15} E_{(0,1,1,1)} - \zeta_{17} E_{(1,1,1,1)} - \zeta_{16} E_{(0,1,1,1,1)} + \zeta_{18} E_{(0,1,1,2,1)} \\ &\quad + \zeta_{19} E_{(1,1,1,1,1)} - \zeta_{21} E_{(1,1,1,2,1)} + \zeta_{22} E_{(1,1,2,2,1)} - \zeta_{27} E_{(1,2,2,3,2,1)} \\ &\quad - 2\zeta_8 c + \frac{\zeta_8}{3}(\alpha_1 + 3\alpha_2 + 2\alpha_3 + 3\alpha_4 + \alpha_5 - \alpha_6) \end{aligned} \tag{5.77}$$

by (2.10), (2.11), (3.7) and (3.46),

$$\begin{aligned} T_9 &= -[\iota(E_{-\alpha_4}), T_7] = \zeta_9(D-8) - \chi\partial_{x_{19}} - \zeta_1 E'_{(1,1,1,2,1)} + \zeta_2 E'_{(0,1,1,2,1)} \\ &\quad + \zeta_4 E'_{(0,1,0,1,1)} - \zeta_6 E'_{(0,0,0,1,1)} + \zeta_5 E'_{(0,1,0,1)} - \zeta_7 E_{-\alpha_4} - \zeta_{11} E_{\alpha_3} + \zeta_{12} E_{\alpha_6} \\ &\quad + \zeta_{13} E_{(0,0,0,0,1,1)} - \zeta_{14} E_{(1,0,1)} + \zeta_{18} E_{(0,0,1,1,1,1)} - \zeta_{20} E_{(0,1,1,1,1,1)} \\ &\quad - \zeta_{21} E_{(1,0,1,1,1,1)} + \zeta_{23} E_{(1,1,1,1,1,1)} + \zeta_{25} E_{(1,1,2,2,1,1)} + \zeta_{26} E_{(1,1,2,2,2,1)} \\ &\quad - 2\zeta_9 c + \frac{\zeta_9}{3}(\alpha_1 + 2\alpha_3 + \alpha_5 + 2\alpha_6) \end{aligned} \tag{5.78}$$

by (2.10), (2.11), (3.5) and (3.44),

$$\begin{aligned} T_{10} &= -[\iota(E_{-\alpha_2}), T_8] = \zeta_{10}(D-8) - \chi\partial_{x_{18}} - \zeta_1 E'_{(1,1,1,1,1,1)} + \zeta_2 E'_{(0,1,1,1,1,1)} \\ &\quad - \zeta_3 E'_{(0,1,0,1,1,1)} + \zeta_6 E'_{(0,0,0,0,1,1)} - \zeta_7 E_{-\alpha_6} - \zeta_8 E_{-\alpha_2} + \zeta_{12} E_{\alpha_4} \\ &\quad - \zeta_{13} E_{(0,0,0,1,1)} - \zeta_{15} E_{(0,0,1,1)} + \zeta_{17} E_{(1,0,1,1)} + \zeta_{16} E_{(0,0,1,1,1)} - \zeta_{20} E_{(0,1,1,2,1)} \\ &\quad - \zeta_{19} E_{(1,0,1,1,1)} + \zeta_{23} E_{(1,1,1,2,1)} - \zeta_{24} E_{(1,1,2,2,1)} - \zeta_{27} E_{(1,1,2,3,2,1)} \\ &\quad - 2\zeta_{10} c + \frac{\zeta_{10}}{3}(\alpha_1 + 2\alpha_3 + 3\alpha_4 + \alpha_5 - \alpha_6) \end{aligned} \tag{5.79}$$

by (2.10), (2.11), (3.3) and (3.42),

$$\begin{aligned} T_{11} &= [\iota(E_{-\alpha_3}), T_9] = \zeta_{11}(D-8) - \chi\partial_{x_{17}} + \zeta_1 E'_{(1,1,2,2,1)} - \zeta_3 E'_{(0,1,1,2,1)} \\ &\quad - \zeta_4 E'_{(0,1,1,1,1)} + \zeta_6 E'_{(0,0,1,1,1)} - \zeta_5 E'_{(0,1,1,1)} + \zeta_7 E'_{(0,0,1,1)} + \zeta_9 E_{-\alpha_3} \end{aligned}$$

$$
\begin{aligned}
&+\zeta_{14}E_{\alpha_1}+\zeta_{15}E_{\alpha_6}+\zeta_{16}E_{(0,0,0,0,1,1)}+\zeta_{18}E_{(0,0,0,1,1,1)}-\zeta_{20}E_{(0,1,0,1,1,1)}\\
&-\zeta_{22}E_{(1,0,1,1,1,1)}+\zeta_{24}E_{(1,1,1,1,1,1)}+\zeta_{25}E_{(1,1,1,2,1,1)}+\zeta_{26}E_{(1,1,1,2,2,1)}\\
&-2\zeta_{11}c+\frac{\zeta_{11}}{3}(\alpha_1-\alpha_3+\alpha_5+2\alpha_6) \qquad (5.80)
\end{aligned}
$$

by (2.10), (2.11), (3.4) and (3.43),

$$
\begin{aligned}
T_{12} \;=\;& -[\iota(E_{-\alpha_4}),T_{10}]=\zeta_{12}(D-8)-\chi\partial_{x_{16}}+\zeta_1E'_{(1,1,1,2,1,1)}-\zeta_2E'_{(0,1,1,2,1,1)}\\
&-\zeta_4E'_{(0,1,0,1,1,1)}+\zeta_6E'_{(0,0,0,1,1,1)}+\zeta_8E'_{(0,1,0,1)}-\zeta_9E_{-\alpha_6}-\zeta_{10}E_{-\alpha_4}\\
&+\zeta_{13}E_{\alpha_5}-\zeta_{15}E_{\alpha_3}+\zeta_{17}E_{(1,0,1)}+\zeta_{18}E_{(0,0,1,1,1)}-\zeta_{20}E_{(0,1,1,1,1)}\\
&-\zeta_{21}E_{(1,0,1,1,1)}+\zeta_{23}E_{(1,1,1,1,1)}+\zeta_{25}E_{(1,1,2,2,1)}-\zeta_{27}E_{(1,1,2,2,2,1)}\\
&-2\zeta_{12}c+\frac{\zeta_{12}}{3}(\alpha_1+2\alpha_3+\alpha_5-\alpha_6) \qquad (5.81)
\end{aligned}
$$

by (2.10), (2.11), (3.5) and (3.44),

$$
\begin{aligned}
T_{13} \;=\;& -[\iota(E_{-\alpha_5}),T_{12}]=\zeta_{13}(D-8)-\chi\partial_{x_{15}}-\zeta_1E'_{(1,1,1,2,2,1)}-\zeta_2E'_{(0,1,1,2,2,1)}\\
&-\zeta_5E'_{(0,1,0,1,1,1)}+\zeta_7E'_{(0,0,0,1,1,1)}-\zeta_8E'_{(0,1,0,1,1)}-\zeta_9E'_{(0,0,0,0,1,1)}+\zeta_{10}E'_{(0,0,0,1,1)}\\
&-\zeta_{12}E_{-\alpha_5}-\zeta_{16}E_{\alpha_3}+\zeta_{19}E_{(1,0,1)}+\zeta_{18}E_{(0,0,1,1)}-\zeta_{20}E_{(0,1,1,1)}\\
&-\zeta_{21}E_{(1,0,1,1)}+\zeta_{23}E_{(1,1,1,1)}-\zeta_{26}E_{(1,1,2,2,1)}-\zeta_{27}E_{(1,1,2,2,1,1)}\\
&-2\zeta_{13}c+\frac{\zeta_{13}}{3}(\alpha_1+2\alpha_3-2\alpha_5-\alpha_6) \qquad (5.82)
\end{aligned}
$$

by (2.10), (2.11), (3.6) and (3.45),

$$
\begin{aligned}
T_{14} \;=\;& -[\iota(E_{-\alpha_1}),T_{11}]=\zeta_{14}(D-8)+\chi\partial_{x_{14}}+\zeta_2E'_{(1,1,2,2,1)}-\zeta_3E'_{(1,1,1,2,1)}\\
&-\zeta_4E'_{(1,1,1,1,1)}+\zeta_6E'_{(1,0,1,1,1)}-\zeta_5E'_{(1,1,1,1)}+\zeta_7E'_{(1,0,1,1)}+\zeta_9E'_{(1,0,1)}\\
&-\zeta_{11}E_{-\alpha_1}-\zeta_{17}E_{\alpha_6}-\zeta_{19}E_{(0,0,0,0,1,1)}-\zeta_{21}E_{(0,0,0,1,1,1)}+\zeta_{23}E_{(0,1,0,1,1,1)}\\
&+\zeta_{22}E_{(0,0,1,1,1,1)}-\zeta_{24}E_{(0,1,1,1,1,1)}-\zeta_{25}E_{(0,1,1,2,1,1)}-\zeta_{26}E_{(0,1,1,2,2,1)}\\
&-2\zeta_{14}c+\frac{\zeta_{14}}{3}(-2\alpha_1-\alpha_3+\alpha_5+2\alpha_6) \qquad (5.83)
\end{aligned}
$$

by (2.10), (2.11), (3.2) and (3.41),

$$
\begin{aligned}
T_{15} \;=\;& [\iota(E_{-\alpha_3}),T_{12}]=\zeta_{15}(D-8)-\chi\partial_{x_{13}}-\zeta_1E'_{(1,1,2,2,1,1)}+\zeta_3E'_{(0,1,1,2,1,1)}\\
&+\zeta_4E'_{(0,1,1,1,1,1)}-\zeta_6E'_{(0,0,1,1,1,1)}-\zeta_8E'_{(0,1,1,1)}+\zeta_{10}E'_{(0,0,1,1)}-\zeta_{11}E_{-\alpha_6}\\
&+\zeta_{12}E_{-\alpha_3}+\zeta_{16}E_{\alpha_5}-\zeta_{17}E_{\alpha_1}+\zeta_{18}E_{(0,0,0,1,1)}-\zeta_{20}E_{(0,1,0,1,1)}\\
&-\zeta_{22}E_{(1,0,1,1,1)}+\zeta_{24}E_{(1,1,1,1,1)}+\zeta_{25}E_{(1,1,1,2,1)}-\zeta_{27}E_{(1,1,1,2,2,1)}\\
&-2\zeta_{15}c+\frac{\zeta_{15}}{3}(\alpha_1-\alpha_3+\alpha_5-\alpha_6) \qquad (5.84)
\end{aligned}
$$

by (2.10), (2.11), (3.4) and (3.43),

$$
\begin{aligned}
T_{16} \;=\;& [\iota(E_{-\alpha_3}),T_{13}]=\zeta_{16}(D-8)-\chi\partial_{x_{12}}+\zeta_1E'_{(1,1,2,2,2,1)}+\zeta_3E'_{(0,1,1,2,2,1)}\\
&+\zeta_5E'_{(0,1,1,1,1,1)}-\zeta_7E'_{(0,0,1,1,1,1)}+\zeta_8E'_{(0,1,1,1,1)}-\zeta_{10}E'_{(0,0,1,1,1)}-\zeta_{11}E'_{(0,0,0,0,1,1)}\\
&+\zeta_{13}E_{-\alpha_3}-\zeta_{15}E_{-\alpha_5}-\zeta_{19}E_{\alpha_1}+\zeta_{18}E_{\alpha_4}-\zeta_{20}E_{(0,1,0,1)}\\
&-\zeta_{22}E_{(1,0,1,1)}+\zeta_{24}E_{(1,1,1,1)}-\zeta_{26}E_{(1,1,1,2,1)}-\zeta_{27}E_{(1,1,1,2,1,1)}\\
&-2\zeta_{16}c+\frac{\zeta_{16}}{3}(\alpha_1-\alpha_3-2\alpha_5-\alpha_6) \qquad (5.85)
\end{aligned}
$$

by (2.10), (2.11), (3.4) and (3.43),

$$\begin{aligned}
T_{17} &= [\iota(E_{-\alpha_1}),T_{15}] = \zeta_{17}(D-8) - \chi\partial_{x_{11}} + \zeta_2 E'_{(1,1,2,2,1,1)} - \zeta_3 E'_{(1,1,1,2,1,1)} \\
&\quad -\zeta_4 E'_{(1,1,1,1,1,1)} + \zeta_6 E'_{(1,0,1,1,1,1)} + \zeta_8 E'_{(1,1,1,1)} - \zeta_{10} E'_{(1,0,1,1)} - \zeta_{12} E'_{(1,0,1)} \\
&\quad +\zeta_{14} E_{-\alpha_6} + \zeta_{15} E_{-\alpha_1} + \zeta_{19} E_{\alpha_5} + \zeta_{21} E_{(0,0,0,1,1)} - \zeta_{23} E_{(0,1,0,1,1)} \\
&\quad -\zeta_{22} E_{(0,0,1,1,1)} + \zeta_{24} E_{(0,1,1,1,1)} + \zeta_{25} E_{(0,1,1,2,1)} - \zeta_{27} E_{(0,1,1,2,2,1)} \\
&\quad -2\zeta_{17} c - \frac{\zeta_{17}}{3}(2\alpha_1 + \alpha_3 - \alpha_5 + \alpha_6)
\end{aligned} \tag{5.86}$$

by (2.10), (2.11), (3.2) and (3.41),

$$\begin{aligned}
T_{18} &= -[\iota(E_{-\alpha_4}),T_{16}] = \zeta_{18}(D-8) - \chi\partial_{x_{10}} - \zeta_1 E'_{(1,1,2,3,2,1)} + \zeta_4 E'_{(0,1,1,2,2,1)} \\
&\quad -\zeta_5 E'_{(0,1,1,2,1,1)} - \zeta_8 E'_{(0,1,1,2,1)} - \zeta_9 E'_{(0,0,1,1,1,1)} - \zeta_{11} E'_{(0,0,0,1,1,1)} - \zeta_{12} E'_{(0,0,1,1,1)} \\
&\quad -\zeta_{13} E'_{(0,0,1,1)} - \zeta_{15} E'_{(0,0,0,1,1)} - \zeta_{16} E_{-\alpha_4} - \zeta_{20} E_{\alpha_2} - \zeta_{21} E_{\alpha_1} \\
&\quad -\zeta_{22} E_{(1,0,1)} - \zeta_{25} E_{(1,1,1,1)} - \zeta_{26} E_{(1,1,1,1,1)} - \zeta_{27} E_{(1,1,1,1,1,1)} \\
&\quad -2\zeta_{16} c + \frac{\zeta_{18}}{3}(\alpha_1 - \alpha_3 - 3\alpha_4 - 2\alpha_5 - \alpha_6)
\end{aligned} \tag{5.87}$$

by (2.10), (2.11), (3.5) and (3.44),

$$\begin{aligned}
T_{19} &= [\iota(E_{-\alpha_1}),T_{16}] = \zeta_{19}(D-8) - \chi\partial_{x_9} - \zeta_2 E'_{(1,1,2,2,2,1)} - \zeta_3 E'_{(1,1,1,2,2,1)} \\
&\quad -\zeta_5 E'_{(1,1,1,1,1,1)} + \zeta_7 E'_{(1,0,1,1,1,1)} - \zeta_8 E'_{(1,1,1,1,1)} + \zeta_{10} E'_{(1,0,1,1,1)} - \zeta_{13} E'_{(1,0,1)} \\
&\quad +\zeta_{14} E'_{(0,0,0,0,1,1)} + \zeta_{16} E_{-\alpha_1} - \zeta_{17} E_{-\alpha_5} + \zeta_{21} E_{\alpha_4} - \zeta_{23} E_{(0,1,0,1)} \\
&\quad -\zeta_{22} E_{(0,0,1,1)} + \zeta_{24} E_{(0,1,1,1)} - \zeta_{26} E_{(0,1,1,2,1)} - \zeta_{27} E_{(0,1,1,2,1,1)} \\
&\quad -2\zeta_{19} c - \frac{\zeta_{19}}{3}(2\alpha_1 + \alpha_3 + 2\alpha_5 + \alpha_6)
\end{aligned} \tag{5.88}$$

by (2.10), (2.11), (3.2) and (3.41),

$$\begin{aligned}
T_{20} &= [\iota(E_{-\alpha_2}),T_{18}] = \zeta_{20}(D-8) - \chi\partial_{x_8} - \zeta_1 E'_{(1,2,2,3,2,1)} - \zeta_6 E'_{(0,1,1,2,2,1)} \\
&\quad +\zeta_7 E'_{(0,1,1,2,1,1)} + \zeta_{10} E'_{(0,1,1,2,1)} + \zeta_9 E'_{(0,1,1,1,1,1)} + \zeta_{11} E'_{(0,1,0,1,1,1)} + \zeta_{12} E'_{(0,1,1,1,1)} \\
&\quad +\zeta_{13} E'_{(0,1,1,1)} + \zeta_{15} E'_{(0,1,0,1,1)} + \zeta_{16} E'_{(0,1,0,1)} + \zeta_{18} E_{-\alpha_2} - \zeta_{23} E_{\alpha_1} \\
&\quad -\zeta_{24} E_{(1,0,1)} - \zeta_{25} E_{(1,0,1,1)} - \zeta_{26} E_{(1,0,1,1,1)} - \zeta_{27} E_{(1,0,1,1,1,1)} \\
&\quad -2\zeta_{20} c + \frac{\zeta_{20}}{3}(\alpha_1 - 3\alpha_2 - \alpha_3 - 3\alpha_4 - 2\alpha_5 - \alpha_6)
\end{aligned} \tag{5.89}$$

by (2.10), (2.11), (3.3) and (3.42),

$$\begin{aligned}
T_{21} &= -[\iota(E_{-\alpha_4}),T_{19}] = \zeta_{21}(D-8) - \chi\partial_{x_7} + \zeta_2 E'_{(1,1,2,3,2,1)} - \zeta_4 E'_{(1,1,1,2,2,1)} \\
&\quad +\zeta_5 E'_{(1,1,1,2,1,1)} + \zeta_8 E'_{(1,1,1,2,1)} + \zeta_9 E'_{(1,0,1,1,1,1)} + \zeta_{12} E'_{(1,0,1,1,1)} + \zeta_{13} E'_{(1,0,1,1)} \\
&\quad +\zeta_{14} E'_{(0,0,0,1,1,1)} - \zeta_{17} E'_{(0,0,0,1,1)} + \zeta_{18} E_{-\alpha_1} - \zeta_{19} E_{-\alpha_4} - \zeta_{22} E_{\alpha_3} \\
&\quad -\zeta_{23} E_{\alpha_2} - \zeta_{25} E_{(0,1,1,1)} - \zeta_{26} E_{(0,1,1,1,1)} - \zeta_{27} E_{(0,1,1,1,1,1)} \\
&\quad -2\zeta_{21} c - \frac{\zeta_{21}}{3}(2\alpha_1 + \alpha_3 + 3\alpha_4 + 2\alpha_5 + \alpha_6)
\end{aligned} \tag{5.90}$$

by (2.10), (2.11), (3.5) and (3.44),

$$\begin{aligned}
T_{22} &= [\iota(E_{-\alpha_3}),T_{21}] = \zeta_{22}(D-8) - \chi\partial_{x_6} - \zeta_3 E'_{(1,1,2,3,2,1)} + \zeta_4 E'_{(1,1,2,2,2,1)} \\
&\quad -\zeta_5 E'_{(1,1,2,2,1,1)} - \zeta_8 E'_{(1,1,2,2,1)} + \zeta_{11} E'_{(1,0,1,1,1,1)} + \zeta_{15} E'_{(1,0,1,1,1)} + \zeta_{16} E'_{(1,0,1,1)}
\end{aligned}$$

$$\begin{aligned}
&-\zeta_{14}E'_{(0,0,1,1,1,1)}+\zeta_{17}E'_{(0,0,1,1,1)}+\zeta_{18}E'_{(1,0,1)}+\zeta_{19}E'_{(0,0,1,1)}+\zeta_{21}E_{-\alpha_3}\\
&-\zeta_{24}E_{\alpha_2}-\zeta_{25}E_{(0,1,0,1)}-\zeta_{26}E_{(0,1,0,1,1)}-\zeta_{27}E_{(0,1,0,1,1,1)}\\
&-2\zeta_{22}c-\frac{\zeta_{22}}{3}(2\alpha_1+4\alpha_3+3\alpha_4+2\alpha_5+\alpha_6) \qquad (5.91)
\end{aligned}$$

by (2.10), (2.11), (3.4) and (3.43),

$$\begin{aligned}
T_{23} \;=\; &[\iota(E_{-\alpha_2}),T_{21}]=\zeta_{23}(D-8)-\chi\partial_{x_5}+\zeta_2E'_{(1,2,2,3,2,1)}+\zeta_6E'_{(1,1,1,2,2,1)}\\
&-\zeta_7E'_{(1,1,1,2,1,1)}-\zeta_{10}E'_{(1,1,1,2,1)}-\zeta_9E'_{(1,1,1,1,1,1)}-\zeta_{12}E'_{(1,1,1,1,1)}-\zeta_{13}E'_{(1,1,1,1)}\\
&-\zeta_{14}E'_{(0,1,0,1,1,1)}+\zeta_{17}E'_{(0,1,0,1,1)}+\zeta_{19}E'_{(0,1,0,1)}+\zeta_{20}E_{-\alpha_1}+\zeta_{21}E_{-\alpha_2}\\
&-\zeta_{24}E_{\alpha_3}-\zeta_{25}E_{(0,0,1,1)}-\zeta_{26}E_{(0,0,1,1,1)}-\zeta_{27}E_{(0,0,1,1,1,1)}\\
&-2\zeta_{23}c-\frac{\zeta_{23}}{3}(2\alpha_1+3\alpha_2+\alpha_3+3\alpha_4+2\alpha_5+\alpha_6) \qquad (5.92)
\end{aligned}$$

by (2.10), (2.11), (3.3) and (3.42),

$$\begin{aligned}
T_{24} \;=\; &[\iota(E_{-\alpha_3}),T_{23}]=\zeta_{24}(D-8)-\chi\partial_{x_4}-\zeta_3E'_{(1,2,2,3,2,1)}-\zeta_6E'_{(1,1,2,2,2,1)}\\
&+\zeta_7E'_{(1,1,2,2,1,1)}+\zeta_{10}E'_{(1,1,2,2,1)}-\zeta_{11}E'_{(1,1,1,1,1,1)}+\zeta_{14}E'_{(0,1,1,1,1,1)}-\zeta_{15}E'_{(1,1,1,1,1)}\\
&-\zeta_{16}E'_{(1,1,1,1)}-\zeta_{17}E'_{(0,1,1,1,1)}-\zeta_{19}E'_{(0,1,1,1)}+\zeta_{20}E'_{(1,0,1)}+\zeta_{22}E_{-\alpha_2}\\
&+\zeta_{23}E_{-\alpha_3}-\zeta_{25}E_{\alpha_4}-\zeta_{26}E_{(0,0,0,1,1)}-\zeta_{27}E_{(0,0,0,1,1,1)}\\
&-2\zeta_{24}c-\frac{\zeta_{24}}{3}(2\alpha_1+3\alpha_2+4\alpha_3+3\alpha_4+2\alpha_5+\alpha_6) \qquad (5.93)
\end{aligned}$$

by (2.10), (2.11), (3.4) and (3.43),

$$\begin{aligned}
T_{25} \;=\; &[\iota(E_{-\alpha_4}),T_{24}]=\zeta_{25}(D-8)-\chi\partial_{x_3}+\zeta_4E'_{(1,2,2,3,2,1)}-\zeta_6E'_{(1,1,2,3,2,1)}\\
&-\zeta_9E'_{(1,1,2,2,1,1)}-\zeta_{11}E'_{(1,1,1,2,1,1)}-\zeta_{12}E'_{(1,1,2,2,1)}+\zeta_{14}E'_{(0,1,1,2,1,1)}-\zeta_{15}E'_{(1,1,1,2,1)}\\
&+\zeta_{18}E'_{(1,1,1,1)}-\zeta_{17}E'_{(0,1,1,2,1)}+\zeta_{20}E'_{(1,0,1,1)}+\zeta_{21}E'_{(0,1,1,1)}+\zeta_{22}E'_{(0,1,0,1)}\\
&+\zeta_{23}E'_{(0,0,1,1)}+\zeta_{24}E_{-\alpha_4}-\zeta_{26}E_{\alpha_5}-\zeta_{27}E_{(0,0,0,0,1,1)}\\
&-2\zeta_{25}c-\frac{\zeta_{25}}{3}(2\alpha_1+3\alpha_2+4\alpha_3+6\alpha_4+2\alpha_5+\alpha_6) \qquad (5.94)
\end{aligned}$$

by (2.10), (2.11), (3.5) and (3.44),

$$\begin{aligned}
T_{26} \;=\; &[\iota(E_{-\alpha_5}),T_{25}]=\zeta_{26}(D-8)-\chi\partial_{x_2}-\zeta_5E'_{(1,2,2,3,2,1)}+\zeta_7E'_{(1,1,2,3,2,1)}\\
&-\zeta_9E'_{(1,1,2,2,2,1)}-\zeta_{11}E'_{(1,1,1,2,2,1)}+\zeta_{13}E'_{(1,1,2,2,1)}+\zeta_{14}E'_{(0,1,1,2,2,1)}\\
&+\zeta_{16}E'_{(1,1,1,2,1)}+\zeta_{18}E'_{(1,1,1,1,1)}+\zeta_{19}E'_{(0,1,1,2,1)}+\zeta_{20}E'_{(1,0,1,1,1)}\\
&+\zeta_{21}E'_{(0,1,1,1,1)}+\zeta_{22}E'_{(0,1,0,1,1)}+\zeta_{23}E'_{(0,0,1,1,1)}+\zeta_{24}E'_{(0,0,0,1,1)}+\zeta_{25}E_{-\alpha_5}\\
&-\zeta_{27}E_{\alpha_6}-2\zeta_{26}c-\frac{\zeta_{26}}{3}(2\alpha_1+3\alpha_2+4\alpha_3+6\alpha_4+5\alpha_5+\alpha_6) \qquad (5.95)
\end{aligned}$$

by (2.10), (2.11), (3.6) and (3.45),

$$\begin{aligned}
T_{27} \;=\; &[\iota(E_{-\alpha_6}),T_{26}]=\zeta_{27}(D-8)-\chi\partial_{x_1}+\zeta_8E'_{(1,2,2,3,2,1)}-\zeta_{10}E'_{(1,1,2,3,2,1)}\\
&+\zeta_{12}E'_{(1,1,2,2,2,1)}+\zeta_{13}E'_{(1,1,2,2,1,1)}+\zeta_{15}E'_{(1,1,1,2,2,1)}+\zeta_{17}E'_{(0,1,1,2,2,1)}\\
&+\zeta_{16}E'_{(1,1,1,2,1,1)}+\zeta_{18}E'_{(1,1,1,1,1,1)}+\zeta_{19}E'_{(0,1,1,2,1,1)}+\zeta_{20}E'_{(1,0,1,1,1,1)}\\
&+\zeta_{21}E'_{(0,1,1,1,1,1)}+\zeta_{22}E'_{(0,1,0,1,1,1)}+\zeta_{23}E'_{(0,0,1,1,1,1)}+\zeta_{24}E'_{(0,0,0,1,1,1)}+\zeta_{26}E_{-\alpha_6}\\
&+\zeta_{25}E'_{(0,0,0,0,1,1)}-2\zeta_{27}c-\frac{\zeta_{27}}{3}(2\alpha_1+3\alpha_2+4\alpha_3+6\alpha_4+5\alpha_5+4\alpha_6) \qquad (5.96)
\end{aligned}$$

by (2.10), (2.11), (3.7) and (3.46). Then $\mathcal{T}=\sum_{r=1}^{27}\mathbb{C}T_r$ forms the 27-dimensional $\mathcal{G}^{E_6}$ of highest weight λ_1. Indeed, the map $\eta_r\mapsto T_r$ for $r\in\overline{1,27}$ determines a $\mathcal{G}^{E_6}$-module isomorphism from U to $\mathcal{T}$ (cf. (3.34)).

Denote

$$T_r'=T_r-\zeta_r(D-2c-8)+\chi\partial_{x_{28-r}}\qquad\text{for }\;r\in\overline{1,27}.\tag{5.97}$$

Easily see that $\mathcal{T}'=\sum_{r=1}^{27}\mathbb{C}T_r'$ forms the 27-dimensional $\mathcal{G}^{E_6}$-module of highest weight λ_1. So we have the $\mathcal{G}^{E_6}$-module isomorphism from $U=\sum_{r=1}^{27}\mathbb{C}\zeta_r$ to $\mathcal{T}'$ determined by $\zeta_r\mapsto T_r'$ for $r\in\overline{1,27}$. The weight set of U is

$$\Pi(U)=\{\sum_{s=1}^{6}b_{r,s}\lambda_s\mid r\in\overline{1,27}\}\tag{5.98}$$

(cf. (5.9)-(5.14) and Table 2 in Section 3). Let $\lambda\in\Lambda^+$. Denote

$$\Upsilon'(\lambda)=\{\lambda+\mu\mid\mu\in\Pi(U),\;\;\lambda+\mu\in\Lambda^+\}\tag{5.99}$$

(cf. (5.27)). Take $M=V(\lambda)$, the irreducible $\mathcal{G}^{E_6}$-module of highest weight λ. It is known that

$$UV(\lambda)=U\otimes_{\mathbb{C}}V(\lambda)\cong\bigoplus_{\lambda'\in\Upsilon'(\lambda)}V(\lambda').\tag{5.100}$$

Given $\lambda'\in\Upsilon'(\lambda)$, we pick a singular vector

$$u=\sum_{r=1}^{27}\zeta_ru_r\tag{5.101}$$

of weight λ' in $UV(\lambda)$, where $u_r\in V(\lambda)$. Moreover, any singular vector of weight λ' in $UV(\lambda)$ is a scalar multiple of u. Note that the vector

$$w=\sum_{r=1}^{27}T_r'(u_r)\tag{5.102}$$

is also a singular vector of weight λ' if it is not zero. Thus

$$w=\flat_{\lambda'}u,\qquad\flat_{\lambda'}\in\mathbb{C}.\tag{5.103}$$

Set

$$\flat(\lambda)=\min\{\flat_{\lambda'}\mid\lambda'\in\Upsilon'(\lambda)\}.\tag{5.104}$$

Theorem 5.6. *The $\mathcal{G}^{E_7}$-module $\widehat{V(\lambda)}$ is irreducible if*

$$c\in\mathbb{C}\setminus\{-8+5\mathbb{N}/3,(1/2)(\flat(\lambda)+\mathbb{N})-4,\ell_\omega(\lambda)+\mathbb{N}\}.\tag{5.105}$$

Proof. Recall that the $\mathcal{G}^{E_7}$-submodule $U(\mathcal{G}_-)(V(\lambda))$ is irreducible by Proposition 4.2. It is enough to prove $\widehat{V(\lambda)}=U(\mathcal{G}_-)(V(\lambda))$. It is obvious that

$$\widehat{V(\lambda)}_{\langle 0\rangle}=V(\lambda)=(U(\mathcal{G}_-)(V(\lambda)))_{\langle 0\rangle}\tag{5.106}$$

(cf. (5.3) and (5.4) with $M = V(\lambda)$). Moreover, Lemma 5.1 with $M = V(\lambda)$, (5.37) and (5.105) imply that $\varphi|_{\widehat{V(\lambda)}_{\langle 1\rangle}}$ is invertible, equivalently,

$$\widehat{V(\lambda)}_{\langle 1\rangle} = (U(\mathcal{G}_-)(V(\lambda)))_{\langle 1\rangle}. \tag{5.107}$$

Suppose that

$$\widehat{V(\lambda)}_{\langle i\rangle} = (U(\mathcal{G}_-)(V(\lambda)))_{\langle i\rangle} \tag{5.108}$$

for $i \in \overline{0,k}$ with $1 \leq k \in \mathbb{N}$.

For any $v \in V(\lambda)$ and $\alpha \in \mathbb{N}^{16}$ such that $|\alpha| = k-1$, we have

$$T_r(x^\alpha v) = x^\alpha[(|\alpha| - 2c - 8)\zeta_r + T_r'](v) + \chi\partial_{x_{28-r}}(x^\alpha)v \in (U(\mathcal{G}_-)(V(\lambda)))_{\langle k+1\rangle} \tag{5.109}$$

for $r \in \overline{1,27}$ by (5.97), (5.108) with $i = k-1, k$. If $k = 1$, then $\alpha = 0$. So $\chi\partial_{x_{28-r}}(x^\alpha)v = 0$. When $k > 1$,

$$\partial_{x_{28-r}}(x^\alpha)v \in \widehat{V(\lambda)}_{\langle k-2\rangle} = (U(\mathcal{G}_-)(V(\lambda)))_{\langle k-2\rangle} \tag{5.110}$$

and so

$$\begin{aligned} & [(\sum_{14\neq p\in\overline{1,26}} \iota(\eta_p)\zeta_{28-p} - \iota(\eta_{14})\zeta_{14} - \iota(\eta_{27})\zeta_1](\partial_{x_{28-r}}(x^\alpha)v) \\ = \ & \chi(24 - 5(|\alpha| - 1) + 3c)(\partial_{x_{28-r}}(x^\alpha)v) \in (U(\mathcal{G}_-)(V(\lambda)))_{\langle k+1\rangle} \end{aligned} \tag{5.111}$$

by Lemma 5.5. Thus (5.105) gives

$$\chi\partial_{x_{28-r}}(x^\alpha)v \in (U(\mathcal{G}_-)(V(\lambda)))_{\langle k+1\rangle} \qquad \text{for } r \in \overline{1,27}. \tag{5.112}$$

Hence in any case,

$$T_r(x^\alpha v) = x^\alpha[(|\alpha| - 2c - 8)\zeta_r + T_r'](v) \in (U(\mathcal{G}_-)(V(\lambda)))_{\langle k+1\rangle} \qquad \text{for } r \in \overline{1,27}. \tag{5.113}$$

On the other hand,

$$V' = \operatorname{Span}\{[(|\alpha| - 2c - 8)\zeta_r + T_r'](v) \mid r \in \overline{1,27},\ v \in V(\lambda)\} \tag{5.114}$$

forms a $\mathcal{G}^{E_6}$-submodule of $UV(\lambda)$ with respect to the action in (5.1). Let u be a $\mathcal{G}^{E_6}$-singular vector in (5.101). Then

$$V' \ni \sum_{r=1}^{27}[(|\alpha| - 2c - 8)\zeta_r + T_r'](u_r) = (|\alpha| - 2c - 8)u + w = (|\alpha| - 2c - 8 + \flat_{\lambda'})u \tag{5.115}$$

by (5.102) and (5.103). Moreover, (5.104) and (5.105) yield $u \in V'$. Since $UV(\lambda)$ is a $\mathcal{G}^{E_6}$-module generated by all the singular vectors, we have $V' = UV(\lambda)$. So

$$x^\alpha UV(\lambda) \subset (U(\mathcal{G}_-)(V(\lambda)))_{\langle k+1\rangle}. \tag{5.116}$$

The arbitrariness of α implies

$$\zeta_r\widehat{V(\lambda)}_{\langle k-1\rangle} \subset (U(\mathcal{G}_-)(V(\lambda)))_{\langle k+1\rangle} \qquad \text{for } r \in \overline{1,27}. \tag{5.117}$$

Given any $f \in \mathcal{A}_k$ and $v \in V(\lambda)$, we have

$$\zeta_r\partial_{x_s}(f)v \in \zeta_r\widehat{V(\lambda)}_{\langle k-1\rangle} \subset (U(\mathcal{G}_-)(V(\lambda)))_{\langle k+1\rangle} \qquad \text{for } r, s \in \overline{1,27}. \tag{5.118}$$

Moreover,

$$\begin{aligned}\eta_s(fv) &= \iota(\eta_s)(fv) = P_s(fv) + f(\tilde{\omega} - c)(x_s v)\\ &\equiv f(k + \tilde{\omega} - c)(x_s v) \ \ (\mathrm{mod} \sum_{r=1}^{27} \zeta_r \widehat{V(\lambda)}_{\langle k-1 \rangle})\end{aligned} \tag{5.119}$$

for $s \in \overline{1,27}$ by (3.58), (3.77)-(3.102), (4.13)-(4.39) and Lemma 5.1. According to (5.37), (5.105), (5.115) and (5.117), we get

$$x_s f v \in (U(\mathcal{G}_-)(V(\lambda)))_{\langle k+1 \rangle} \qquad \text{for } \ s \in \overline{1,27}. \tag{5.120}$$

Thus (5.108) holds for $i = k+1$. By induction on k, (5.108) holds for any $i \in \mathbb{N}$, that is, $\widehat{V(\lambda)} = U(\mathcal{G}_-)(V(\lambda))$. $\qquad\Box$

When $\lambda = 0$, $V(0)$ is the one-dimensional trivial module and $\ell_\omega(0) = \flat(0) = 0$. So we have:

Corollary 6.7. *The* $\mathcal{G}^{E_7}$*-module* $\widehat{V(0)}$ *is irreducible if* $c \in \mathbb{C} \setminus \{5\mathbb{N}/3 - 8, \mathbb{N}/2 - 4\}$.

Next we consider $\lambda = k\lambda_1$. In this case,

$$\Upsilon(\lambda) = \{k\lambda_1 + \lambda_6, (k-1)\lambda_1 + \lambda_2, (k-1)\lambda_1\} \tag{5.121}$$

by (5.32) and Tables 1, 2. Thus we have

$$\ell_\omega(\lambda) = -16 - \frac{4k}{3} \tag{5.122}$$

by (5.37). Moreover,

$$\Upsilon'(\lambda) = \{(k+1)\lambda_1, (k-1)\lambda_1 + \lambda_3, (k-1)\lambda_1 + \lambda_6\} \tag{5.123}$$

by Table 2 and (5.99).

We define a representation of $\mathcal{G}^{E_6}$ on $\mathcal{C} = \mathbb{C}[z_1, ..., z_{27}]$ determined via (3.35)-(3.47) with U replaced by $\mathcal{C}$ and ζ_i replaced by z_i. In fact,

$$E_{\alpha_1}|_{\mathcal{C}} = z_1\partial_{z_2} + z_{11}\partial_{z_{14}} - z_{15}\partial_{z_{17}} - z_{16}\partial_{z_{19}} - z_{18}\partial_{z_{21}} - z_{20}\partial_{z_{23}}, \tag{5.124}$$

$$E_{\alpha_2}|_{\mathcal{C}} = z_4\partial_{z_6} + z_5\partial_{z_7} + z_8\partial_{z_{10}} - z_{18}\partial_{z_{20}} - z_{21}\partial_{z_{23}} - z_{22}\partial_{z_{24}}, \tag{5.125}$$

$$E_{\alpha_3}|_{\mathcal{C}} = z_2\partial_{z_3} - z_9\partial_{z_{11}} - z_{12}\partial_{z_{15}} - z_{13}\partial_{z_{16}} - z_{21}\partial_{z_{22}} - z_{23}\partial_{z_{24}}, \tag{5.126}$$

$$E_{\alpha_4}|_{\mathcal{C}} = z_3\partial_{z_4} + z_7\partial_{z_9} + z_{10}\partial_{z_{12}} + z_{16}\partial_{z_{18}} + z_{19}\partial_{z_{21}} - z_{24}\partial_{z_{25}}, \tag{5.127}$$

$$E_{\alpha_5}|_{\mathcal{C}} = z_4\partial_{z_5} + z_6\partial_{z_7} + z_{12}\partial_{z_{13}} + z_{15}\partial_{z_{16}} + z_{17}\partial_{z_{19}} - z_{25}\partial_{z_{26}}, \tag{5.128}$$

$$E_{\alpha_6}|_{\mathcal{C}} = z_5\partial_{z_8} + z_7\partial_{z_{10}} + z_9\partial_{z_{12}} + z_{11}\partial_{z_{15}} - z_{14}\partial_{z_{17}} - z_{26}\partial_{z_{27}}, \tag{5.129}$$

$$E_{(1,0,1)}|_{\mathcal{C}} = -z_1\partial_{z_3} - z_9\partial_{z_{14}} + z_{12}\partial_{z_{17}} + z_{13}\partial_{z_{19}} - z_{18}\partial_{z_{22}} - z_{20}\partial_{z_{24}}, \tag{5.130}$$

$$E_{(0,1,0,1)}|_{\mathcal{C}} = z_3\partial_{z_6} - z_5\partial_{z_9} - z_8\partial_{z_{12}} - z_{16}\partial_{z_{20}} - z_{19}\partial_{z_{23}} - z_{22}\partial_{z_{25}}, \tag{5.131}$$

$$E_{(0,0,1,1)}|_{\mathcal{C}} = -z_2\partial_{z_4} - z_7\partial_{z_{11}} - z_{10}\partial_{z_{15}} + z_{13}\partial_{z_{18}} - z_{19}\partial_{z_{22}} - z_{23}\partial_{z_{25}}, \tag{5.132}$$

$$E_{(0,0,0,1,1)}|_{\mathcal{C}} = -z_3\partial_{z_5} + z_6\partial_{z_9} - z_{10}\partial_{z_{13}} + z_{15}\partial_{z_{18}} + z_{17}\partial_{z_{21}} - z_{24}\partial_{z_{26}}, \tag{5.133}$$

$$E_{(0,0,0,0,1,1)}|_{\mathcal{C}} = -z_4\partial_{z_8} - z_6\partial_{z_{10}} + z_9\partial_{z_{13}} + z_{11}\partial_{z_{16}} - z_{14}\partial_{z_{19}} + z_{25}\partial_{z_{27}}, \tag{5.134}$$

$$E_{(1,0,1,1)}|_{\mathcal{C}} = z_1\partial_{z_4} - z_7\partial_{z_{14}} + z_{10}\partial_{z_{17}} - z_{13}\partial_{z_{21}} - z_{16}\partial_{z_{22}} - z_{20}\partial_{z_{25}}, \tag{5.135}$$

$$E_{(0,1,1,1)}|_{\mathcal{C}} = -z_2\partial_{z_6} + z_5\partial_{z_{11}} + z_8\partial_{z_{15}} - z_{13}\partial_{z_{20}} + z_{19}\partial_{z_{z_{24}}} - z_{21}\partial_{z_{25}}, \tag{5.136}$$

$$E_{(0,1,0,1,1)}|_{\mathcal{C}} = -z_3\partial_{z_7} - z_4\partial_{z_9} + z_8\partial_{z_{13}} - z_{15}\partial_{z_{20}} - z_{17}\partial_{z_{23}} - z_{22}\partial_{z_{26}}, \tag{5.137}$$

$$E_{(0,0,1,1,1)}|_{\mathcal{C}} = z_2\partial_{z_5} - z_6\partial_{z_{11}} + z_{10}\partial_{z_{16}} + z_{12}\partial_{z_{18}} - z_{17}\partial_{z_{22}} - z_{23}\partial_{z_{26}}, \tag{5.138}$$

$$E_{(0,0,0,1,1,1)}|_{\mathcal{A}} = z_3\partial_{z_8} - z_6\partial_{z_{12}} - z_7\partial_{z_{13}} + z_{11}\partial_{z_{18}} - z_{14}\partial_{z_{21}} + z_{24}\partial_{z_{27}}, \tag{5.139}$$

$$E_{(1,1,1,1)}|_{\mathcal{C}} = z_1\partial_{z_6} + z_5\partial_{z_{14}} - z_8\partial_{z_{17}} + z_{13}\partial_{z_{23}} + z_{16}\partial_{z_{24}} - z_{18}\partial_{z_{25}}, \tag{5.140}$$

$$E_{(1,0,1,1,1)}|_{\mathcal{C}} = -z_1\partial_{z_5} - z_6\partial_{z_{14}} - z_{10}\partial_{z_{19}} - z_{12}\partial_{z_{21}} - z_{15}\partial_{z_{22}} - z_{20}\partial_{z_{26}}, \tag{5.141}$$

$$E_{(0,1,1,1,1)}|_{\mathcal{C}} = z_2\partial_{z_7} + z_4\partial_{z_{11}} - z_8\partial_{z_{16}} - z_{12}\partial_{z_{20}} + z_{17}\partial_{z_{24}} - z_{21}\partial_{z_{26}}, \tag{5.142}$$

$$E_{(0,1,0,1,1,1)}|_{\mathcal{C}} = z_3\partial_{z_{10}} + z_4\partial_{z_{12}} + z_5\partial_{z_{15}} - z_{11}\partial_{z_{20}} + z_{14}\partial_{z_{23}} + z_{22}\partial_{z_{27}}, \tag{5.143}$$

$$E_{(0,0,1,1,1,1)}|_{\mathcal{C}} = -z_2\partial_{z_8} + z_6\partial_{z_{15}} + z_7\partial_{z_{16}} + z_9\partial_{z_{18}} + z_{14}\partial_{z_{22}} + z_{23}\partial_{z_{27}}, \tag{5.144}$$

$$E_{(1,1,1,1,1}|_{\mathcal{C}} = -z_1\partial_{z_7} + z_4\partial_{z_{14}} + z_8\partial_{z_{19}} + z_{12}\partial_{z_{23}} + z_{15}\partial_{z_{24}} - z_{18}\partial_{z_{26}}, \tag{5.145}$$

$$E_{(1,0,1,1,1,1)}|_{\mathcal{C}} = z_1\partial_{z_8} - z_6\partial_{z_{17}} - z_7\partial_{z_{19}} - z_9\partial_{z_{21}} - z_{11}\partial_{z_{22}} + z_{20}\partial_{z_{27}}, \tag{5.146}$$

$$E_{(0,1,1,2,1)}|_{\mathcal{C}} = -z_2\partial_{z_9} + z_3\partial_{z_{11}} + z_8\partial_{z_{18}} - z_{10}\partial_{z_{20}} + z_{17}\partial_{z_{25}} + z_{19}\partial_{z_{26}}, \tag{5.147}$$

$$E_{(0,1,1,1,1,1)}|_{\mathcal{C}} = -z_2\partial_{z_{10}} - z_4\partial_{z_{15}} - z_5\partial_{z_{16}} - z_9\partial_{z_{20}} - z_{14}\partial_{z_{24}} + z_{21}\partial_{z_{27}}, \tag{5.148}$$

$$E_{(1,1,1,2,1)}|_{\mathcal{C}} = z_1\partial_{z_9} + z_3\partial_{z_{14}} - z_8\partial_{z_{21}} + z_{10}\partial_{z_{23}} + z_{15}\partial_{z_{25}} + z_{16}\partial_{z_{26}}, \tag{5.149}$$

$$E_{(1,1,1,1,1,1)}|_{\mathcal{C}} = z_1\partial_{z_{10}} + z_4\partial_{z_{17}} + z_5\partial_{z_{19}} + z_9\partial_{z_{23}} + z_{11}\partial_{z_{24}} + z_{18}\partial_{z_{27}}, \tag{5.150}$$

$$E_{(0,1,1,2,1,1)}|_{\mathcal{C}} = z_2\partial_{z_{12}} - z_3\partial_{z_{15}} + z_5\partial_{z_{18}} - z_7\partial_{z_{20}} - z_{14}\partial_{z_{25}} + z_{19}\partial_{z_{27}}, \tag{5.151}$$

$$E_{(1,1,2,2,1)}|_{\mathcal{C}} = -z_1\partial_{z_{11}} - z_2\partial_{z_{14}} + z_8\partial_{z_{22}} - z_{10}\partial_{z_{24}} + z_{12}\partial_{z_{25}} + z_{13}\partial_{z_{26}}, \tag{5.152}$$

$$E_{(1,1,1,2,1,1)}|_{\mathcal{C}} = -z_1\partial_{z_{12}} + z_3\partial_{z_{17}} - z_5\partial_{z_{21}} + z_7\partial_{z_{23}} + z_{11}\partial_{z_{25}} + z_{16}\partial_{z_{27}}, \tag{5.153}$$

$$E_{(0,1,1,2,2,1)}|_{\mathcal{C}} = -z_2\partial_{z_{13}} + z_3\partial_{z_{16}} + z_4\partial_{z_{18}} - z_6\partial_{z_{20}} - z_{14}\partial_{z_{26}} + z_{17}\partial_{z_{27}}, \tag{5.154}$$

$$E_{(1,1,2,2,1,1)}|_{\mathcal{C}} = z_1\partial_{z_{15}} - z_2\partial_{z_{17}} + z_5\partial_{z_{22}} - z_7\partial_{z_{24}} + z_9\partial_{z_{25}} + z_{13}\partial_{z_{27}}, \tag{5.155}$$

$$E_{(1,1,1,2,2,1)}|_{\mathcal{C}} = z_1\partial_{z_{13}} - z_3\partial_{z_{19}} - z_4\partial_{z_{21}} + z_6\partial_{z_{23}} + z_{11}\partial_{z_{26}} + z_{15}\partial_{z_{27}}, \tag{5.156}$$

$$E_{(1,1,2,2,2,1)}|_{\mathcal{C}} = -z_1\partial_{z_{16}} + z_2\partial_{z_{19}} + z_4\partial_{z_{22}} - z_6\partial_{z_{24}} + z_9\partial_{z_{26}} + z_{12}\partial_{z_{27}}, \tag{5.157}$$

$$E_{(1,1,2,3,2,1)}|_{\mathcal{C}} = z_1\partial_{z_{18}} - z_2\partial_{z_{21}} + z_3\partial_{z_{22}} - z_6\partial_{z_{25}} + z_7\partial_{z_{26}} + z_{10}\partial_{z_{27}}, \tag{5.158}$$

$$E_{(1,2,2,3,2,1)}|_{\mathcal{C}} = z_1\partial_{z_{20}} - z_2\partial_{z_{23}} + z_3\partial_{z_{24}} - z_4\partial_{z_{25}} + z_5\partial_{z_{26}} + z_8\partial_{z_{27}}, \tag{5.159}$$

$$\alpha_r|_{\mathcal{C}} = \sum_{i=1}^{27} b_{i,r} z_i \partial_{z_i} \qquad \text{for } \ r \in \overline{1,6} \tag{5.160}$$

with $b_{i,r}$ given Table 2,

$$E_{-\alpha}|_{\mathcal{C}} = -\tau'(E_\alpha) \qquad \text{for } \ \alpha\imath\Phi_{E_6}, \tag{5.161}$$

where τ' is a linear transformation on the differential operators in $z_1, ..., z_{27}$ such that

$$\tau'(z_i\partial_{z_j}) = z_j\partial_{z_i} \qquad \text{for } \ i,j \in \overline{1,27}. \tag{5.162}$$

Then the $\mathcal{G}^{E_6}$-submodule $\mathcal{N}_k$ generated by z_1^k is isomorphic to $V(k\lambda_1)$.

We calculate:

$$z_1^{k-1}z_2=-\frac{1}{k}E_{-\alpha_1}(z_1^k)\in\mathcal{N}_k,\quad z_1^{k-1}z_3=-E_{-\alpha_3}(z_1^{k-1}z_2)\in\mathcal{N}_k, \tag{5.163}$$

$$z_1^{k-1}z_4=-E_{-\alpha_4}(z_1^{k-1}z_3)\in\mathcal{N}_k,\quad z_1^{k-1}z_5=-E_{-\alpha_5}(z_1^{k-1}z_4)\in\mathcal{N}_k, \tag{5.164}$$

$$z_1^{k-1}z_6=-E_{-\alpha_2}(z_1^{k-1}z_4)\in\mathcal{N}_k,\quad z_1^{k-1}z_7=-E_{-\alpha_2}(z_1^{k-1}z_5)\in\mathcal{N}_k, \tag{5.165}$$

$$z_1^{k-1}z_9=-E_{-\alpha_4}(z_1^{k-1}z_7)\in\mathcal{N}_k,\quad z_1^{k-1}z_{11}=E_{-\alpha_3}(z_1^{k-1}z_9)\in\mathcal{N}_k, \tag{5.166}$$

$$-E_{-\alpha_1}(z_1^{k-1}z_{11})=z_1^{k-1}z_{14}+(k-1)z_1^{k-2}z_2z_{11}\in\mathcal{N}_k, \tag{5.167}$$

$$-E_{-\alpha_3}(z_1^{k-1}z_{14}+(k-1)z_1^{k-2}z_2z_{11})=(k-1)z_3z_{11}\in\mathcal{N}_k, \tag{5.168}$$

$$E_{\alpha_3}((k-1)z_3z_{11})=(k-1)(z_2z_{11}-z_3z_9)\in\mathcal{N}_k, \tag{5.169}$$

$$E_{\alpha_4}E_{-\alpha_4}[(k-1)(z_2z_{11}-z_3z_9)]=(k-1)(z_3z_9+z_4z_7)\in\mathcal{N}_k, \tag{5.170}$$

$$-E_{\alpha_5}E_{-\alpha_5}[(k-1)(z_3z_9+z_4z_7)]=(k-1)(z_4z_7+z_5z_6)\in\mathcal{N}_k. \tag{5.171}$$

Now we take $M=\mathcal{N}_k$ in our earlier settings. First $\zeta_1z_1^k$ is a singular vector of weight $(k+1)\lambda_1$ in $U\mathcal{N}_k$. By (5.69), (5.160) and Table 2,

$$T_1'(z_1^k)=\frac{\zeta_1}{3}(4\alpha_1+3\alpha_2+5\alpha_3+6\alpha_4+4\alpha_5+2\alpha_6)(z_1^k)=\frac{4k}{3}\zeta_1z_1^k. \tag{5.172}$$

So $\flat_{(k+1)\lambda_1}=4k/3$. Next $\zeta_1z_1^{k-1}z_2-\zeta_2z_1^k$ is a singular vector of weight $(k-1)\lambda_1+\lambda_3$ in $U\mathcal{N}_k$. According to (5.69), (5.71), (5.160) and Table 2:

$$\begin{aligned}&T_1'(z_1^{k-1}z_2)-T_2'(z_1^k)\\ =\ &\zeta_2z_1^k+\frac{4k-3}{3}\zeta_1z_1^{k-1}z_2-k\zeta_1z_1^{k-1}z_2-\frac{k}{3}\zeta_2z_1^k=\frac{k-3}{3}(\zeta_1z_1^{k-1}z_2-\zeta_2z_1^k),\end{aligned} \tag{5.173}$$

which gives $\flat_{(k-1)\lambda_1+\lambda_3}=k/3-1$.

Expressions (3.35)-(3.40), (5.124)-(5.129) and (5.163)-(5.171) show that

$$\begin{aligned}u\ =\ &\zeta_1[4z_1^{k-1}z_{14}+(k-1)z_1^{k-2}(z_2z_{11}+z_3z_9-z_4z_7+z_5z_6)]-(k+3)z_1^{k-1}\\ &\times[\zeta_2z_{11}+\zeta_3z_9-\zeta_4z_7+\zeta_5z_6+\zeta_6z_5-\zeta_7z_4+\zeta_9z_3+\zeta_{11}z_2-\zeta_{14}z_1]\end{aligned} \tag{5.174}$$

is a singular vector of weight $(k-1)\lambda_1+\lambda_6$ in $U\mathcal{N}_k$. We find

$$\begin{aligned}&T_1'[4z_1^{k-1}z_{14}+(k-1)z_1^{k-2}(z_2z_{11}+z_3z_9-z_4z_7+z_5z_6)]\\ &-(k+3)[T_2'(z_1^{k-1}z_{11})+T_3'(z_1^{k-1}z_9)-T_4'(z_1^{k-1}z_7)+T_5'(z_1^{k-1}z_6)\\ &+T_6'(z_1^{k-1}z_5)-T_7'(z_1^{k-1}z_4)+T_9'(z_1^{k-1}z_3)+T_{11}'(z_1^{k-1}z_2)-T_{14}'(z_1^k)]\\ =\ &(k+3)z_1^{k-1}(\zeta_2z_{11}+\zeta_3z_9-\zeta_4z_7+\zeta_5z_6+\zeta_6z_5-\zeta_7z_4+\zeta_9z_3+\zeta_{11}z_2)\\ &+(4k/3-2)\zeta_1[4z_1^{k-1}z_{14}+(k-1)z_1^{k-2}(z_2z_{11}+z_3z_9-z_4z_7+z_5z_6)]\\ &-(k+3)z_1^{k-2}[2z_1(\zeta_1z_{14}-\zeta_3z_9+\zeta_4z_7-\zeta_5z_6-\zeta_6z_5+\zeta_7z_4-\zeta_9z_3+\zeta_{14}z_1)\\ &+2z_1(\zeta_1z_{14}-\zeta_2z_{11}+\zeta_4z_7-\zeta_5z_6-\zeta_6z_5+\zeta_7z_4-\zeta_{11}z_2+\zeta_{14}z_1)\\ &+2z_1(\zeta_1z_{14}-\zeta_2z_{11}-\zeta_3z_9-\zeta_5z_6-\zeta_6z_5-\zeta_9z_3-\zeta_{11}z_2+\zeta_{14}z_1)\\ &+2z_1(\zeta_1z_{14}-\zeta_2z_{11}-\zeta_3z_9+\zeta_4z_7+\zeta_7z_4-\zeta_9z_3-\zeta_{11}z_2+\zeta_{14}z_1)\\ &+2(k-1)\zeta_1(z_2z_{11}+z_3z_9-z_4z_7+z_5z_6)-(2k/3+1)z_1(\zeta_2z_{11}+\zeta_3z_9\\ &-\zeta_4z_7+\zeta_5z_6+\zeta_6z_5-\zeta_7z_4+\zeta_9z_3+\zeta_{11}z_2)+(2k/3)\zeta_{14}z_1^2]\end{aligned}$$

$$
\begin{aligned}
&= (k+3)z_1^{k-1}(\zeta_2 z_{11}+\zeta_3 z_9-\zeta_4 z_7+\zeta_5 z_6+\zeta_6 z_5-\zeta_7 z_4+\zeta_9 z_3+\zeta_{11} z_2)\\
&\quad +(4k/3-2)\zeta_1[4z_1^{k-1}z_{14}+(k-1)z_1^{k-2}(z_2 z_{11}+z_3 z_9-z_4 z_7+z_5 z_6)]-(k+3)z_1^{k-2}\\
&\quad \times[2(4\zeta_1 z_1 z_{14}+(k-1)\zeta_1(z_2 z_{11}+z_3 z_9-z_4 z_7+z_5 z_6))+(2k/3+8)\zeta_{14} z_1^2\\
&\quad -(2k/3+7)z_1(\zeta_2 z_{11}+\zeta_3 z_9-\zeta_4 z_7+\zeta_5 z_6+\zeta_6 z_5-\zeta_7 z_4+\zeta_9 z_3+\zeta_{11} z_2)]\\
&= -(2k/3+8)u
\end{aligned}
\tag{5.175}
$$

by (5.69), (5.71)-(5.96), (5.124)-(5.161) and Table 2. Thus $\flat_{(k-1)\lambda_1+\lambda_6} = -(2k/3+8)$. Therefore,

$$\flat(k\lambda_1) = -(2k/3+8). \tag{5.176}$$

Corollary 5.8. *The $\mathcal{G}^{E_7}$-module $\widehat{V(k\lambda_1)}$ is irreducible if*

$$c \in \mathbb{C} \setminus \{-8+5\mathbb{N}/3, \mathbb{N}/2-8-k/3, \mathbb{N}-16-4k/3\}. \tag{5.177}$$

References

[**A**] J. Adams, *Lectures on Exceptional Lie Groups*, The University of Chicago Press Ltd., London, 1996.

[**BKR**] L. Brink, S. Kim and P. Ramond, $E_{7(7)}$ on the light cone, *J. High Energy Phys.* (2008), no. 6, 034, 18pp.

[**Br1**] R. Brown, A minimal representation for the Lie algebra of type E_7, *Illiois J. Math.* **12** (1968), 190-200.

[**Br2**] R. Brown, Groups of type E_7, *J. Reine Angew. Math.* **236** (1969), 79-102.

[**BK**] R. Brylinski and B. Kostant, Minimal representations of E_6, E_7, and E_8 and the generalized Capelli identity, *Proc. Nar. Acad. Sci. U.S.A.* **91**, no. 7., 2469-2472.

[**Bl**] L. Borsten, $E_{7(7)}$ invariant measures of entanglement, *Fortschr. Phys.* **56** (2008), no. 7-9, 842-848.

[**CY**] Y. Choi and S. Yoon, Homology of the double and the triple loop spaces of E_7, E_7 and E_8, *Manuscripta Math.* **103** (2000), 101-116.

[**Cv**] V. Chernousov, The kernel of the Rost invariant, Serre's conjecture II and the Hasse principle for quasi-split groups D_4, E_6, E_7, *Math. Ann.* **326** (2003), no. 2, 297-330.

[**Cb**] B. Cooperstein, The fifty-six-dimensional module for E_7.I. A four form for E_7, *J. Algebra* **173** (1995), 361-389.

[**Co**] O. Cvitanović, Negative dimensions and E_7 symmetry, *Nuclear Phys. B* **188** (1981), no. 2, 373-396.

[**DFT**] R. D'Auria, S. Ferrara and M. Trigiante, $E_{7(7)}$ symmetry and dual gauge algebras of M-theory on a twisted even-torus, *Nuclear Phys. B* **732** (2006), 389-400.

[Dt] T. De Medts, A characterization of quadratic forms of type E_6, E_7 and E_8, *J. Algebra* **252** (2002), no. 2, 394-410.

[Dl] L. Dickson, A class of groups in an arbitrary realm connected with the configuration of the 27 lines on a cubic surface, *J. Math.* **33** (1901), 145-123.

[Dd1] D. Doković, Explicit Cayley triples in real forms of E_7, *Pacific J. Math.* **191** (1999), 1-23.

[Dd2] D. Doković, The closure daigram for nilpotent orbits of the split real form of E_7, *Reprent. Theory* **5** (2001), 284-316.

[DF] M. Duff and S. Ferrara, E_7 and the tripartite entanglement of seven qubits, *Phys. Rev. D* **76** (2007), no. 2, 025018, 7pp.

[Fau] J. Faulkner, A geometry for E_7, *Trans. Amer. Math. Soc.* **167** (1972), 49-58.

[FFP] N. Fernádez, W. Garcia Fuerres and A. Perelomov, Quantum trogonometric Calogero-Sutherland model, irreducible characters and Clebsch-Gordan sries for the exceptional algebra E_7, *J. Math. Phys.* **46** (2005), no. 10, 103505, 27pp.

[Fer] J. Ferrar, On the classification of Freudenthal triple systems and Lie algebras of type E_7, *J. Algebra* **62** (1980), 276-282.

[Gr] R. Garibaldi, Structurable algebras and groups of type E_6 and E_7, *J. Algebra* **236** (2001), no. 2, 651-691.

[Gd] D. Ginzburg, On standard L-functions for E_6 and E_7, *J. Reine Angew. Math.* **465** (1995), 101-131.

[GR] R. Griess and A. Ryba, Embeddings of $U_3(8)$, $Sz(8)$ and the Rudvalis group in algebraic groups of type E_7, *Invent. Math.* **116** (1994), 215-141.

[HKKT] S. Han, J. Kim, I. Koh and Y. Tanii, Supersymmetrization of the six-dimensional anomaly-free $E_6 \times E_7 \times U(1)$ theory with Lorentz Chern-Simons term, *Phys. Lett. B* **177** (1986), no. 2, 167-170.

[H] J. E. Humphreys, *Introduction to Lie Algebras and Representation Theory*, Springer-Verlag New York Inc., 1972.

[K] V. Kac, *Infinite Dimensional Lie Algebras,* Third Edition, Cambridge University Press, 1990.

[KK] A. Kato and Y. Kitazawa, E_7-type modular invariant Wess-Zumino theory and Gepner's string compactification, *Nuclear Phys. B* **319** (1989), no. 2, 474-490.

[KR] P. Kleidman and A. Ryba, Kostant's conjecture holds for $E_7 : L_2(37) < E_7(C)$, *J. Algebra* **161** (1993), 535-540.

[KMR] P. Kleidman, U. Meierfrankenfeld and A. Ryba, $HS < E_7(5)$, *J. London Math. Soc. (2)* **60** (1999), 95-107.

[KLN] A. Kono, J. Lin and O. Nishimura, Characterization of the mod 3 cohomology of E_7, *Proc. Amer. Math. Soc.* **131** (2003), no. 10, 3289-3295.

[M] Z. Ma, The spectrum-dependent solutions of the Yang-Baxter equation for quantum E_6 and E_7, *J. Phys. A* **23** (1990), no. 23, 5513-5522.

[P] E. Plotkin, On the stability of the K_1-functor for Chevalley groups of type E_7, *J. Algebra* **210** (1998), 67-85.

[R] P. Ramond, Is there an exceptional group in your future? E_7 and the travails of the symmetry breaking, *Nuclear Phys. B* **126** (1977), no. 3, 509-524.

[Sj] J. Sekiguchi, Configurations of seven lines on the real projective plane and the root system of type E_7, *J. Math. Soc. Japan* **51** (1999), no. 4, 987-1013.

[Se] E. Shult, Embeddings and hyperplanes of the Lie incidence geometry of type $E_{7,1}$, *J. Geom.* **59** (1997), 152-172.

[U] K. Ukai, b-functions of the prehomogeneous vector space arising from a cuspidal character sheaf of E_7, *J. Algebra* **237** (2001), 358-381.

[W] R. Weiss, Moufang quadangles of type E_6 and E_7, *J. Reine Angew. Math.* **590** (2006), 189-226.

[X1] X. Xu, *Kac-Moody Algebras and Their Representations*, China Science Press, 2007.

[X2] X. Xu, A cubic E_6-generalization of the classical theorem on harmonic polynomials, *J. Lie Theory* **21** (2011), 145-164.

[X3] X. Xu, Representations of Lie algebras and coding theory, *J. Lie Theory* **22** (2012), no. 3, 647-682.

[X4] X. Xu, A new functor from D_5-**Mod** to E_6-**Mod**, *arXiv:1112.3792v1[math.RT].*

[XZ] X. Xu and Y. Zhao, Generalized conformal representations of orthogonal Lie algebras, *arXiv:1105.1254v1[math.RT].*

[ZX] Y. Zhao and X. Xu, Generalized projective representations for sl(n+1), *J. Algebra* **328** (2011), 132-154.